

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problems Mailbox.**

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 5 : C12N 15/12, C07K 13/00 C12P 21/08, G01N 33/68 A61K 37/02		A1	(11) International Publication Number: WO 91/18982 (43) International Publication Date: 12 December 1991 (12.12.91)
(21) International Application Number: PCT/US91/03498 (22) International Filing Date: 17 May 1991 (17.05.91) (30) Priority data: 534,193 5 June 1990 (05.06.90) US 573,576 24 August 1990 (24.08.90) US 627,071 13 December 1990 (13.12.90) US (71) Applicant: IMMUNEX CORPORATION [US/US]; 51 University Street, Seattle, WA 98101 (US). (72) Inventors: SIMS, John, E. ; 314 N.E. 82nd Street, Seattle, WA 98115 (US). COSMAN, David, J. ; 9808 Mandus Olson Road N.E., Bainbridge Island, WA 98110 (US). LUPTON, Stephen, D. ; 7323 Sandpoint Way N.E., #211, Seattle, WA 98115 (US). MOSLEY, Bruce, A. ; 11331 - 36th Avenue N.E., Seattle, WA 98125 (US). DOWER, Steven, K. ; 2620 East Lake Sammamish Parkway N.E., Redmond, WA 98052 (US).		(74) Agent: HALLQUIST, Scott, G.; Immunex Corporation, 51 University Street, Seattle, WA 98101 (US). (81) Designated States: AU, CA, FI, KR, NO. Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.	

(54) Title: TYPE II INTERLEUKIN-1 RECEPTORS

510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000

## (57) Abstract

Type II IL-1 receptor (type II IL-1R) proteins, DNAs and expression vectors encoding type II IL-1R, and processes for producing type II IL-1R as products of recombinant cell culture, are disclosed.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MN	Mongolia
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GN	Guinea	NL	Netherlands
BJ	Benin	GR	Greece	NO	Norway
BR	Brazil	HU	Hungary	PL	Poland
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LI	Liechtenstein	SU	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TG	Togo
DE	Germany	MC	Monaco	US	United States of America
DK	Denmark				

## **TITLE**

### **Type II Interleukin-1 Receptors**

## **BACKGROUND OF THE INVENTION**

The present invention relates generally to cytokine receptors, and more specifically, to Interleukin-1 receptors.

Interleukin-1 $\alpha$  (IL-1 $\alpha$ ) and Interleukin-1 $\beta$  and (IL-1 $\beta$ ) are distantly related polypeptide hormones which play a central role in the regulation of immune and inflammatory responses. These two proteins act on a variety of cell types and have multiple biological activities. The diversity of biological activity ascribed to IL-1 $\alpha$  and IL-1 $\beta$  is mediated by specific plasma membrane receptors which bind both IL-1 $\alpha$  and IL-1 $\beta$ . Due to the wide range of biological activities mediated by IL-1 $\alpha$  and IL-1 $\beta$  it was originally believed that the IL-1 receptors should be highly conserved in a variety of species and expressed on a large variety of cells.

Structural characterization by ligand affinity cross-linking techniques has demonstrated that, despite their significant divergence in sequence, IL-1 $\alpha$  and IL-1 $\beta$  bind to the same cell surface receptor molecule on T cells and fibroblasts (Dower et al., *Nature (London)* 324:266, 1986; Bird et al., *Nature (London)* 324:263, 1986; Dower et al., *Proc. Natl. Acad. Sci. USA* 83:1060, 1986). The IL-1 receptor on murine and human T cells has been identified by cDNA expression cloning and N-terminal sequence analysis as an integral membrane glycoprotein that binds IL-1 $\alpha$  and IL-1 $\beta$  and has a molecular weight of 80,000 kDa (Sims et al., *Science* 241:585, 1988; Sims et al., *Proc. Natl. Acad. Sci. USA* 86:8946, 1989).

It is now clear, however, that this 80 kDa IL-1 receptor protein does not mediate all the diverse biological effects of IL-1. Subsequent affinity cross-linking studies indicate that IL-1 receptors on the Epstein Barr virus (EBV)-transformed human B cell lines VDS-O and 3B6, the EBV-positive Burkitt's lymphoma cell line Raji, and the murine pre-B cell line 70Z/3, have a molecular weight of 60,000 to 68,000 kDa (Matsushima et al., *J. Immunol.* 136:4496, 1986; Bensimon et al., *J. Immunol.* 142:2290, 1989; Bensimon et al., *J. Immunol.* 143:1168, 1989; Horuk et al., *J. Biol. Chem.* 262:16275, 1987; Chizzonite et al., *Proc. Natl. Acad. Sci. USA* 86:8029, 1989; Bomsztyk et al., *Proc. Natl. Acad. Sci. USA* 86:8034, 1989). Moreover, comparison of the biochemical properties and kinetic analysis of the IL-1 receptor in the Raji B cell line with EL-4 murine T lymphoma cell line showed that Raji cells had lower binding affinity but much higher receptor density per cell than a subclone of EL-4 T cells (Horuk et al., *J. Biol. Chem.* 262:16275, 1987). Raji cells also failed to internalize IL-1 and demonstrated altered receptor binding affinities

with IL-1 analogs. (Horuk et al., *J. Biol. Chem.* 262:16275, 1987). These data suggest that the IL-1 receptors expressed on B cells (referred to herein as type II IL-1 receptors) are different from IL-1 receptors detected on T cells and other cell types (referred to herein as type I IL-1 receptors).

In order to study the structural and biological characteristics of type II IL-1R and the role played by type II IL-1R in the responses of various cell populations to IL-1 stimulation, or to use type II IL-1R effectively in therapy, diagnosis, or assay, homogeneous compositions are needed. Such compositions are theoretically available via purification of receptors expressed by cultured cells, or by cloning and expression of genes encoding the receptors. Prior to the present invention, however, several obstacles prevented these goals from being achieved.

First, no cell lines have previously been known to express high levels of type II IL-1R constitutively and continuously, and cell lines known to express type II IL-1R did so only in low numbers (500 to 2,000 receptors/cell) which impeded efforts to purify receptors in amounts sufficient for obtaining amino acid sequence information or generating monoclonal antibodies. The low numbers of receptors has also precluded any practical translation assay-based method of cloning.

Second, the significant differences in DNA sequence between type I IL-1R and type II IL-1R has precluded cross-hybridization using a murine type IL-1R cDNA (Bomsztyk et al., *Proc. Natl. Acad. Sci. USA* 86:8034, 1989, and Chizzonite et al., *Proc. Natl. Acad. Sci. USA* 86:8029, 1989).

Third, even if a protein composition of sufficient purity could be obtained to permit N-terminal protein sequencing, the degeneracy of the genetic code may not permit one to define a suitable probe without considerable additional experimentation. Many iterative attempts may be required to define a probe having the requisite specificity to identify a hybridizing sequence in a cDNA library. Although direct expression cloning techniques avoid the need for repetitive screening using different probes of unknown specificity and have been useful in cloning other receptors (e.g., type I IL-1R), they are not sufficiently sensitive to be suitable for using in identifying type II IL-1R clones from cDNA libraries derived from cells expressing low numbers of type II IL-1R.

Thus, efforts to purify the type II IL-1R or to clone or express genes encoding type II IL-1R have been significantly impeded by lack of purified receptor, a suitable source of receptor mRNA, and by a sufficiently sensitive cloning technique.

### **SUMMARY OF THE INVENTION**

The present invention provides purified and homogeneous type II IL-1R proteins and isolated DNA sequences encoding type II IL-1R proteins, in particular, human type II

IL-1R, or analogs thereof. Preferably, such DNA sequences are selected from the group consisting of (a) cDNA clones having a nucleotide sequence derived from the coding region of a native type II IL-1R gene, such as clone 75; (b) DNA sequences capable of hybridization to the cDNA clones of (a) under moderately stringent conditions and which encode biologically active IL-1R molecules; and (c) DNA sequences which are degenerate as a result of the genetic code to the DNA sequences defined in (a) and (b) and which encode biologically active IL-1R molecules. The present invention also provides recombinant expression vectors comprising the DNA sequences defined above, recombinant type II IL-1R molecules produced using the recombinant expression vectors, and processes for producing the recombinant type II IL-1R molecules utilizing the expression vectors.

The present invention also provides substantially purified and homogeneous proteins and protein compositions comprising type II IL-1R.

The present invention also provides compositions for use in therapy, diagnosis, assay of type II IL-1R, or in raising antibodies to type II IL-1R, comprising effective quantities of soluble native or recombinant receptor proteins prepared according to the foregoing processes.

These and other aspects of the present invention will become evident upon reference to the following detailed description.

### **BRIEF DESCRIPTION OF THE DRAWINGS**

FIG. 1 is a schematic diagram of the expression plasmid pDC406. cDNA molecules inserted at the Sal site are transcribed and translated using regulatory elements derived from HIV and adenovirus. pDC406 contains origins of replication derived from SV40, Epstein-Barr virus and pBR322.

FIG. 2 is a schematic diagram of the human and murine type II IL-1 receptors and the various human and murine clones used to determine the sequences. Thin lines represent untranslated regions, while the coding region is depicted by a box. The sections encoding the signal peptide are filled in; the transmembrane regions are cross-hatched; and the cytoplasmic portions are stippled. Potential N-linked glycosylation sites are marked by inverted triangles. The predicted immunoglobulin-like disulfide bonds are also indicated by dashes connecting two sulfide molecules (S—S).

FIG. 3 compares the amino acid sequences of the human and murine type II IL-1 receptors (as deduced from the cDNA clones) with the amino acid sequences of the human and murine type I IL-1 receptors (Sims et al., *Proc. Natl. Acad. Sci. USA* 86:8946, 1989; Sims et al., *Science* 241:585, 1988) and the amino acid sequences of the ST2 cellular gene

(Tominaga, *FEBS Lett.* 258:301, 1989) and the B15R open reading frame of vaccinia virus (Smith and Chan, *J. Gen. Virology* 72:511, 1991). Numbering begins with the initiating methionine. The predicted position of the signal peptide cleavage in each sequences was determined according to the method described by von Heijne, *Nucl. Acids. Res.* 14:4683, 1986, and is indicated by a gap between the putative signal peptide and the main body of the protein. The predicated transmembrane and cytoplasmic regions for the type II IL-1 receptors are shown on the bottom line, and are separated from one another by a gap. Residues conserved in all four IL-1 receptor sequences are presented in white on a black background. Residues conserved in type II receptors that are also found in one of the other sequences are shaded; residues conserved in type I IL-1 receptors that are found in one of the other sequences are boxed. Cysteine residues involved in forming the disulfide bonds characteristic of the immunoglobulin fold are marked with solid dots, while the extra two pairs of cysteines found in the type I IL-1 receptor and in some of the other sequences are indicated by stars. The approximate boundaries of domains 1, 2 and 3 are indicated above the lines. The predicted signal peptide cleavage in the type II IL-1 receptors follow Ala13, resulting in an unusually short signal peptide and an N-terminal extension of 12 (human) or 23 (mouse) amino acids beyond the point corresponding to the mature N-terminus of the human or mouse type I IL-1 receptor. Other less favored but still acceptable sites of cleavage in the murine type II IL-1 receptor are after Thr15 or Pro17. This sequence alignment was made by hand and does not represent an objectively optimized alignment of the sequences. The nucleotide and amino acid sequences of the full length and soluble human and murine type II IL-1 receptor cDNAs are also set forth in the Sequence Listing herein.

FIG. 4 shows an autoradiograph of an SDS/PAGE gel with crosslinked IL-1 receptors. Cells expressing IL-1 receptors were cross-linked to  $^{125}\text{I}$ -IL-1 in the absence or presence of the cognate unlabeled IL-1 competitor, extracted, electrophoresed and autoradiographed as described in Example 6. Recombinant receptors were expressed transiently in CV1/EBNA cells. The cell lines used for cross-linking to natural receptors were KB (ATCC CCL 1717) (for human type I IL-1R), CB23 (for human type II IL-1R), EL4 (ATCC TIB 39) (for murine type I IL-1R), and 70Z/3 (ATCC TIB 158) (for murine type II IL-1R).

## DETAILED DESCRIPTION OF THE INVENTION

### Definitions

"IL-1" refers collectively to IL-1 $\alpha$  and IL-1 $\beta$ .

"Type II Interleukin-1 receptor" and "type II IL-1R" refer to proteins which are capable of binding Interleukin-1 (IL-1) molecules and, in their native configuration as mammalian plasma membrane proteins, play a role in transducing the signal provided by IL-1 to a cell. The mature full-length human type II IL-1R is a glycoprotein having an apparent molecular weight of approximately 60-68 kDa. Specific examples of type II IL-1R proteins are shown in SEQ ID NO:1 and SEQ ID NO:12. As used herein, the above terms include analogs or subunits of native type II IL-1R proteins with IL-1-binding or signal transducing activity. Specifically included are truncated or soluble forms of type II IL-1R protein, as defined below. In the absence of any species designation, type II IL-1R refers generically to mammalian type II IL-1R, which includes, but is not limited to, human, murine, and bovine type II IL-1R. Similarly, in the absence of any specific designation for deletion mutants, the term type II IL-1R means all forms of type II IL-1R, including mutants and analogs which possess type II IL-1R biological activity. "Interleukin-1 Receptor" or "IL-1R" refers collectively to type I IL-1 receptor and type II IL-1 receptor.

"Soluble type II IL-1R" as used in the context of the present invention refer to proteins, or substantially equivalent analogs, which are substantially similar to all or part of the extracellular region of a native type II IL-1R, and are secreted by the cell but retain the ability to bind IL-1 or inhibit IL-1 signal transduction activity via cell surface bound IL-1R proteins. Soluble type II IL-1R proteins may also include part of the transmembrane region, provided that the soluble type II IL-1R protein is capable of being secreted from the cell. Specific examples of soluble type II IL-1R proteins include proteins having the sequence of amino acids 1-330 or amino acids 1-333 of SEQ ID NO:1 and amino acids 1-342 and amino acids 1-345 of SEQ ID NO:12. Inhibition of IL-1 signal transduction activity can be determined using primary cells or cells lines which express an endogenous IL-1R and which are biologically responsive to IL-1 or which, when transfected with recombinant IL-1R DNAs, are biologically responsive to IL-1. The cells are then contacted with IL-1 and the resulting metabolic effects examined. If an effect results which is attributable to the action of the ligand, then the recombinant receptor has signal transduction activity. Exemplary procedures for determining whether a polypeptide has signal transduction activity are disclosed by Idzerda et al., *J. Exp. Med.* 171:861 (1990); Curtis et al., *Proc. Natl. Acad. Sci. USA* 86:3045 (1989); Prywes et al., *EMBO J.* 5:2179 (1986) and Chou et al., *J. Biol. Chem.* 262:1842 (1987).

The term "isolated" or "purified", as used in the context of this specification to define the purity of type II IL-1R protein or protein compositions, means that the protein or protein composition is substantially free of other proteins of natural or endogenous origin and contains less than about 1% by mass of protein contaminants residual of production processes. Such compositions, however, can contain other proteins added as stabilizers, carriers, excipients or co-therapeutics. Type II IL-1R is "isolated" if it is detectable as a single protein band in a polyacrylamide gel by silver staining.

The term "substantially similar," when used to define either amino acid or nucleic acid sequences, means that a particular subject sequence, for example, a mutant sequence, varies from a reference sequence by one or more substitutions, deletions, or additions, the net effect of which is to retain biological activity of the type II IL-1R protein as may be determined, for example, in a type II IL-1R binding assays, such as is described in Example 5 below. Alternatively, nucleic acid subunits and analogs are "substantially similar" to the specific DNA sequences disclosed herein if: (a) the DNA sequence is derived from the coding region of SEQ ID NO:1 or SEQ ID NO:12; (b) the DNA sequence is capable of hybridization to DNA sequences of (a) under moderately stringent conditions (25% formamide, 42°C, 2xSSC) or alternatively under more stringent conditions (50% formamide, 50°C, 2xSSC or 50% formamide, 42°C, 2xSSC) and which encode biologically active IL-1R molecules; or DNA sequences which are degenerate as a result of the genetic code to the DNA sequences defined in (a) or (b) and which encode biologically active IL-1R molecules.

"Recombinant," as used herein, means that a protein is derived from recombinant (e.g., microbial or mammalian) expression systems. "Microbial" refers to recombinant proteins made in bacterial or fungal (e.g., yeast) expression systems. As a product, "recombinant microbial" defines a protein essentially free of native endogenous substances and unaccompanied by associated native glycosylation. Protein expressed in most bacterial cultures, e.g., *E. coli*, will be free of glycan; protein expressed in yeast may have a glycosylation pattern different from that expressed in mammalian cells.

"Biologically active," as used throughout the specification as a characteristic of type II IL-1R, means either that a particular molecule shares sufficient amino acid sequence similarity with SEQ ID NO:2 or SEQ ID NO:13 to be capable of binding detectable quantities of IL-1, preferably at least 0.01 nmoles IL-1 per nanomole type II IL-1R, or, in the alternative, shares sufficient amino acid sequence similarity to be capable of transmitting an IL-1 stimulus to a cell, for example, as a component of a hybrid receptor construct. More preferably, biologically active type II IL-1R within the scope of the present invention is capable of binding greater than 0.1 nanomoles IL-1 per nanomole receptor, and most preferably, greater than 0.5 nanomoles IL-1 per nanomole receptor.

"DNA sequence" refers to a DNA polymer, in the form of a separate fragment or as a component of a larger DNA construct, which has been derived from DNA isolated at least once in substantially pure form, i.e., free of contaminating endogenous materials and in a quantity or concentration enabling identification, manipulation, and recovery of the sequence and its component nucleotide sequences by standard biochemical methods, for example, using a cloning vector. Such sequences are preferably provided in the form of an open reading frame uninterrupted by internal nontranslated sequences, or introns, which are typically present in eukaryotic genes. However, it will be evident that genomic DNA containing the relevant sequences could also be used. Sequences of non-translated DNA may be present 5' or 3' from the open reading frame, where the same do not interfere with manipulation or expression of the coding regions.

"Nucleotide sequence" refers to a heteropolymer of deoxyribonucleotides. DNA sequences encoding the proteins provided by this invention are assembled from cDNA fragments and short oligonucleotide linkers, or from a series of oligonucleotides, to provide a synthetic gene which is capable of being expressed in a recombinant transcriptional unit.

"Recombinant expression vector" refers to a plasmid comprising a transcriptional unit comprising an assembly of (1) a genetic element or elements having a regulatory role in gene expression, for example, promoters or enhancers, (2) a structural or coding sequence which is transcribed into mRNA and translated into protein, and (3) appropriate transcription and translation initiation and termination sequences. Structural elements intended for use in yeast expression systems preferably include a leader sequence enabling extracellular secretion of translated protein by a host cell. Alternatively, where recombinant protein is expressed without a leader or transport sequence, it may include an N-terminal methionine residue. This residue may optionally be subsequently cleaved from the expressed recombinant protein to provide a final product.

"Recombinant microbial expression system" means a substantially homogeneous monoculture of suitable host microorganisms, for example, bacteria such as *E. coli* or yeast such as *S. cerevisiae*, which have stably integrated a recombinant transcriptional unit into chromosomal DNA or carry the recombinant transcriptional unit as a component of a resident plasmid. Generally, cells constituting the system are the progeny of a single ancestral transformant. Recombinant expression systems as defined herein will express heterologous protein upon induction of the regulatory elements linked to the DNA sequence or synthetic gene to be expressed.

### Type II IL-1R Proteins and Analogs

The present invention provides isolated recombinant mammalian type II IL-1R polypeptides. The type II IL-1R proteins of the present invention include, by way of example, primate, human, murine, canine, feline, bovine, ovine, equine, caprine and porcine type II IL-1R. Type II IL-1R can be obtained by cross species hybridization, using a single stranded cDNA derived from the human or murine type II IL-1R DNA sequence, for example, human clone 75, as a hybridization probe to isolate type II IL-1R cDNAs from mammalian cDNA libraries. Like most mammalian genes, mammalian type II IL-1R is presumably encoded by multi-exon genes. Alternative mRNA constructs which can be attributed to different mRNA splicing events following transcription, and which share large regions of identity or similarity with the cDNAs claimed herein, are considered to be within the scope of the present invention. DNA sequences which encode IL-1R-II, possibly in the form of alternate splicing arrangements, can be isolated from the following cells and tissues: B lymphoblastoid lines (such as CB23, CB33, Raji, RPMI1788, ARH77), resting and especially activated peripheral blood T cells, monocytes, the monocytic cell line THP1, neutrophils, bone marrow, placenta, endothelial cells, keratinocytes (especially activated), and HepG2 cells.

Derivatives of type II IL-1R within the scope of the invention also include various structural forms of the primary protein which retain biological activity. Due to the presence of ionizable amino and carboxyl groups, for example, a type II IL-1R protein may be in the form of acidic or basic salts, or may be in neutral form. Individual amino acid residues may also be modified by oxidation or reduction.

The primary amino acid structure may be modified by forming covalent or aggregative conjugates with other chemical moieties, such as glycosyl groups, lipids, phosphate, acetyl groups and the like, or by creating amino acid sequence mutants. Covalent derivatives are prepared by linking particular functional groups to type II IL-1R amino acid side chains or at the N- or C-termini. Other derivatives of type II IL-1R within the scope of this invention include covalent or aggregative conjugates of type II IL-1R or its fragments with other proteins or polypeptides, such as by synthesis in recombinant culture as N-terminal or C-terminal fusions. For example, the conjugated peptide may be a signal (or leader) polypeptide sequence at the N-terminal region of the protein which co-translationally or post-translationally directs transfer of the protein from its site of synthesis to its site of function inside or outside of the cell membrane or wall (e.g., the yeast  $\alpha$ -factor leader). Type II IL-1R protein fusions can comprise peptides added to facilitate purification or identification of type II IL-1R (e.g., poly-His). The amino acid sequence of type II IL-1R can also be linked to the peptide Asp-Tyr-Lys-Asp-Asp-Asp-Lys (DYKDDDDK) (Hopp et al., *BioTechnology* 6:1204,1988.) The latter sequence is highly antigenic and

provides an epitope reversibly bound by a specific monoclonal antibody, enabling rapid assay and facile purification of expressed recombinant protein. This sequence is also specifically cleaved by bovine mucosal enterokinase at the residue immediately following the Asp-Lys pairing. Fusion proteins capped with this peptide may also be resistant to intracellular degradation in *E. coli*.

Type II IL-1R derivatives may also be used as immunogens, reagents in receptor-based immunoassays, or as binding agents for affinity purification procedures of IL-1 or other binding ligands. Type II IL-1R derivatives may also be obtained by cross-linking agents, such as M-maleimidobenzoyl succinimide ester and N-hydroxysuccinimide, at cysteine and lysine residues. Type II IL-1R proteins may also be covalently bound through reactive side groups to various insoluble substrates, such as cyanogen bromide-activated, bisoxirane-activated, carbonyldiimidazole-activated or tosyl-activated agarose structures, or by adsorbing to polyolefin surfaces (with or without glutaraldehyde cross-linking). Once bound to a substrate, type II IL-1R may be used to selectively bind (for purposes of assay or purification) anti-type II IL-1R antibodies or IL-1.

The present invention also includes type II IL-1R with or without associated native-pattern glycosylation. Type II IL-1R expressed in yeast or mammalian expression systems, e.g., COS-7 cells, may be similar or slightly different in molecular weight and glycosylation pattern than the native molecules, depending upon the expression system. Expression of type II IL-1R DNAs in bacteria such as *E. coli* provides non-glycosylated molecules. Functional mutant analogs of mammalian type II IL-1R having inactivated N-glycosylation sites can be produced by oligonucleotide synthesis and ligation or by site-specific mutagenesis techniques. These analog proteins can be produced in a homogeneous, reduced-carbohydrate form in good yield using yeast expression systems. N-glycosylation sites in eukaryotic proteins are characterized by the amino acid triplet Asn-A<sub>1</sub>-Z, where A<sub>1</sub> is any amino acid except Pro, and Z is Ser or Thr. In this sequence, asparagine provides a side chain amino group for covalent attachment of carbohydrate. Examples of N-glycosylation sites in human type II IL-1R are amino acids 66-68, 72-74, 112-114, 219-221, and 277-279 in SEQ ID NO:1. Such sites can be eliminated by substituting another amino acid for Asn or for residue Z, deleting Asn or Z, or inserting a non-Z amino acid between A<sub>1</sub> and Z, or an amino acid other than Asn between Asn and A<sub>1</sub>.

Type II IL-1R derivatives may also be obtained by mutations of type II IL-1R or its subunits. A type II IL-1R mutant, as referred to herein, is a polypeptide homologous to type II IL-1R but which has an amino acid sequence different from native type II IL-1R because of a deletion, insertion or substitution.

Bioequivalent analogs of type II IL-1R proteins may be constructed by, for example, making various substitutions of residues or sequences or deleting terminal or internal residues or sequences not needed for biological activity. For example, cysteine residues can be deleted or replaced with other amino acids to prevent formation of unnecessary or incorrect intramolecular disulfide bridges upon renaturation. Other approaches to mutagenesis involve modification of adjacent dibasic amino acid residues to enhance expression in yeast systems in which KEX2 protease activity is present. Generally, substitutions should be made conservatively; i.e., the most preferred substitute amino acids are those having physiochemical characteristics resembling those of the residue to be replaced. Similarly, when a deletion or insertion strategy is adopted, the potential effect of the deletion or insertion on biological activity should be considered. Substantially similar polypeptide sequences, as defined above, generally comprise a like number of amino acids sequences, although C-terminal truncations for the purpose of constructing soluble type II IL-1Rs will contain fewer amino acid sequences. In order to preserve the biological activity of type II IL-1Rs, deletions and substitutions will preferably result in homologous or conservatively substituted sequences, meaning that a given residue is replaced by a biologically similar residue. Examples of conservative substitutions include substitution of one aliphatic residue for another, such as Ile, Val, Leu, or Ala for one another, or substitutions of one polar residue for another, such as between Lys and Arg; Glu and Asp; or Gln and Asn. Other such conservative substitutions, for example, substitutions of entire regions having similar hydrophobicity characteristics, are well known. Moreover, particular amino acid differences between human, murine and other mammalian type II IL-1Rs is suggestive of additional conservative substitutions that may be made without altering the essential biological characteristics of type II IL-1R.

Subunits of type II IL-1R may be constructed by deleting terminal or internal residues or sequences. The present invention contemplates, for example, C terminal deletions which result in soluble type II IL-1R constructs corresponding to all or part of the extracellular region of type II IL-1R. The resulting protein preferably retains its ability to bind IL-1. Particularly preferred sequences include those in which the transmembrane region and intracellular domain of type II IL-1R are deleted or substituted with hydrophilic residues to facilitate secretion of the receptor into the cell culture medium. Soluble type II IL-1R proteins may also include part of the transmembrane region, provided that the soluble type II IL-1R protein is capable of being secreted from the cell. For example, soluble human type II IL-1R may comprise the sequence of amino acids 1-333 or amino acids 1-330 of SEQ ID NO:1 and amino acids 1-345 and amino acids 1-342 of SEQ ID NO:12. Alternatively, soluble type II IL-1R proteins may be derived by deleting the C-terminal region of a type II IL-1R within the extracellular region which are not necessary

for IL-1 binding. For example, C-terminal deletions may be made to proteins having the sequence of SEQ ID NO:1 and SEQ ID NO:12 following amino acids 313 and 325, respectively. These amino acids are cysteines which are believed to be necessary to maintain the tertiary structure of the type II IL-1R molecule and permit binding of the type II IL-1R molecule to IL-1. Soluble type II IL-1R constructs are constructed by deleting the 3'-terminal region of a DNA encoding the type II IL-1R and then inserting and expressing the DNA in appropriate expression vectors. Exemplary methods of constructing such soluble proteins are described in Examples 2 and 4. The resulting soluble type II IL-1R proteins are then assayed for the ability to bind IL-1, as described in Example 5. Both the DNA sequences encoding such soluble type II IL-1Rs and the biologically active soluble type II IL-1R proteins resulting from such constructions are contemplated to be within the scope of the present invention.

Mutations in nucleotide sequences constructed for expression of analog type II IL-1R must, of course, preserve the reading frame phase of the coding sequences and preferably will not create complementary regions that could hybridize to produce secondary mRNA structures such as loops or hairpins which would adversely affect translation of the receptor mRNA. Although a mutation site may be predetermined, it is not necessary that the nature of the mutation *per se* be predetermined. For example, in order to select for optimum characteristics of mutants at a given site, random mutagenesis may be conducted at the target codon and the expressed type II IL-1R mutants screened for the desired activity.

Not all mutations in the nucleotide sequence which encodes type II IL-1R will be expressed in the final product, for example, nucleotide substitutions may be made to enhance expression, primarily to avoid secondary structure loops in the transcribed mRNA (see EPA 75,444A, incorporated herein by reference), or to provide codons that are more readily translated by the selected host, e.g., the well-known *E. coli* preference codons for *E. coli* expression.

Mutations can be introduced at particular loci by synthesizing oligonucleotides containing a mutant sequence, flanked by restriction sites enabling ligation to fragments of the native sequence. Following ligation, the resulting reconstructed sequence encodes an analog having the desired amino acid insertion, substitution, or deletion.

Alternatively, oligonucleotide-directed site-specific mutagenesis procedures can be employed to provide an altered gene having particular codons altered according to the substitution, deletion, or insertion required. Exemplary methods of making the alterations set forth above are disclosed by Walder et al. (*Gene* 42:133, 1986); Bauer et al. (*Gene* 37:73, 1985); Craik (*BioTechniques*, January 1985, 12-19); Smith et al. (*Genetic Engineering: Principles and Methods*, Plenum Press, 1981); and U.S. Patent Nos.

4,518,584 and 4,737,462 disclose suitable techniques, and are incorporated by reference herein.

Both monovalent forms and polyvalent forms of type II IL-1R are useful in the compositions and methods of this invention. Polyvalent forms possess multiple type II IL-1R binding sites for IL-1 ligand. For example, a bivalent soluble type II IL-1R may consist of two tandem repeats of the extracellular region of type II IL-1R, separated by a linker region. Alternate polyvalent forms may also be constructed, for example, by chemically coupling type II IL-1R to any clinically acceptable carrier molecule, a polymer selected from the group consisting of Ficoll, polyethylene glycol or dextran using conventional coupling techniques. Alternatively, type II IL-1R may be chemically coupled to biotin, and the biotin-type II IL-1R conjugate then allowed to bind to avidin, resulting in tetravalent avidin/biotin/type II IL-1R molecules. Type II IL-1R may also be covalently coupled to dinitrophenol (DNP) or trinitrophenol (TNP) and the resulting conjugate precipitated with anti-DNP or anti-TNP-IgM, to form decameric conjugates with a valency of 10 for type II IL-1R binding sites.

A recombinant chimeric antibody molecule may also be produced having type II IL-1R sequences substituted for the variable domains of either or both of the immunoglobulin molecule heavy and light chains and having unmodified constant region domains. For example, chimeric type II IL-1R/IgG<sub>1</sub> may be produced from two chimeric genes -- a type II IL-1R/human  $\kappa$  light chain chimera (type II IL-1R/C $\kappa$ ) and a type II IL-1R/human  $\gamma$ 1 heavy chain chimera (type II IL-1R/C $\gamma$ 1). Following transcription and translation of the two chimeric genes, the gene products assemble into a single chimeric antibody molecule having type II IL-1R displayed bivalently. Such polyvalent forms of type II IL-1R may have enhanced binding affinity for IL-1 ligand. Additional details relating to the construction of such chimeric antibody molecules are disclosed in WO 89/09622 and EP 315062.

#### Expression of Recombinant Type II IL-1R

The present invention provides recombinant expression vectors to amplify or express DNA encoding type II IL-1R. Recombinant expression vectors are replicable DNA constructs which have synthetic or cDNA-derived DNA fragments encoding mammalian type II IL-1R or bioequivalent analogs operably linked to suitable transcriptional or translational regulatory elements derived from mammalian, microbial, viral or insect genes. A transcriptional unit generally comprises an assembly of (1) a genetic element or elements having a regulatory role in gene expression, for example, transcriptional promoters or enhancers, (2) a structural or coding sequence which is transcribed into mRNA and translated into protein, and (3) appropriate transcription and translation initiation and

termination sequences, as described in detail below. Such regulatory elements may include an operator sequence to control transcription, a sequence encoding suitable mRNA ribosomal binding sites. The ability to replicate in a host, usually conferred by an origin of replication, and a selection gene to facilitate recognition of transformants may additionally be incorporated. DNA regions are operably linked when they are functionally related to each other. For example, DNA for a signal peptide (secretory leader) is operably linked to DNA for a polypeptide if it is expressed as a precursor which participates in the secretion of the polypeptide; a promoter is operably linked to a coding sequence if it controls the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to permit translation. Generally, operably linked means contiguous and, in the case of secretory leaders, contiguous and in reading frame. Structural elements intended for use in yeast expression systems preferably include a leader sequence enabling extracellular secretion of translated protein by a host cell. Alternatively, where recombinant protein is expressed without a leader or transport sequence, it may include an N-terminal methionine residue. This residue may optionally be subsequently cleaved from the expressed recombinant protein to provide a final product.

DNA sequences encoding mammalian type II IL-1Rs which are to be expressed in a microorganism will preferably contain no introns that could prematurely terminate transcription of DNA into mRNA; however, premature termination of transcription may be desirable, for example, where it would result in mutants having advantageous C-terminal truncations, for example, deletion of a transmembrane region to yield a soluble receptor not bound to the cell membrane. Due to code degeneracy, there can be considerable variation in nucleotide sequences encoding the same amino acid sequence. Other embodiments include sequences capable of hybridizing to clone 75 under moderately stringent conditions (50°C, 2x SSC) and other sequences hybridizing or degenerate to those which encode biologically active type II IL-1R polypeptides.

Recombinant type II IL-1R DNA is expressed or amplified in a recombinant expression system comprising a substantially homogeneous monoculture of suitable host microorganisms, for example, bacteria such as *E. coli* or yeast such as *S. cerevisiae*, which have stably integrated (by transformation or transfection) a recombinant transcriptional unit into chromosomal DNA or carry the recombinant transcriptional unit as a component of a resident plasmid. Generally, cells constituting the system are the progeny of a single ancestral transformant. Recombinant expression systems as defined herein will express heterologous protein upon induction of the regulatory elements linked to the DNA sequence or synthetic gene to be expressed.

Transformed host cells are cells which have been transformed or transfected with type II IL-1R vectors constructed using recombinant DNA techniques. Transformed host

cells ordinarily express type II IL-1R, but host cells transformed for purposes of cloning or amplifying type II IL-1R DNA do not need to express type II IL-1R. Expressed type II IL-1R will be deposited in the cell membrane or secreted into the culture supernatant, depending on the type II IL-1R DNA selected. Suitable host cells for expression of mammalian type II IL-1R include prokaryotes, yeast or higher eukaryotic cells under the control of appropriate promoters. Prokaryotes include gram negative or gram positive organisms, for example *E. coli* or bacilli. Higher eukaryotic cells include established cell lines of mammalian origin as described below. Cell-free translation systems could also be employed to produce mammalian type II IL-1R using RNAs derived from the DNA constructs of the present invention. Appropriate cloning and expression vectors for use with bacterial, fungal, yeast, and mammalian cellular hosts are described by Pouwels et al. (*Cloning Vectors: A Laboratory Manual*, Elsevier, New York, 1985), the relevant disclosure of which is hereby incorporated by reference.

Prokaryotic expression hosts may be used for expression of type II IL-1R that do not require extensive proteolytic and disulfide processing. Prokaryotic expression vectors generally comprise one or more phenotypic selectable markers, for example a gene encoding proteins conferring antibiotic resistance or supplying an autotrophic requirement, and an origin of replication recognized by the host to ensure amplification within the host. Suitable prokaryotic hosts for transformation include *E. coli*, *Bacillus subtilis*, *Salmonella typhimurium*, and various species within the genera *Pseudomonas*, *Streptomyces*, and *Staphylococcus*, although others may also be employed as a matter of choice.

Useful expression vectors for bacterial use can comprise a selectable marker and bacterial origin of replication derived from commercially available plasmids comprising genetic elements of the well known cloning vector pBR322 (ATCC 37017). Such commercial vectors include, for example, pKK223-3 (Pharmacia Fine Chemicals, Uppsala, Sweden) and pGEM1 (Promega Biotec, Madison, WI, USA). These pBR322 "backbone" sections are combined with an appropriate promoter and the structural sequence to be expressed. *E. coli* is typically transformed using derivatives of pBR322, a plasmid derived from an *E. coli* species (Bolivar et al., *Gene* 2:95, 1977). pBR322 contains genes for ampicillin and tetracycline resistance and thus provides simple means for identifying transformed cells.

Promoters commonly used in recombinant microbial expression vectors include the  $\beta$ -lactamase (penicillinase) and lactose promoter system (Chang et al., *Nature* 275:615, 1978; and Goeddel et al., *Nature* 281:544, 1979), the tryptophan (trp) promoter system (Goeddel et al., *Nucl. Acids Res.* 8:4057, 1980; and EPA 36,776) and tac promoter (Maniatis, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory, p. 412, 1982). A particularly useful bacterial expression system employs the phage  $\lambda$  P<sub>L</sub>

promoter and cI857ts thermolabile repressor. Plasmid vectors available from the American Type Culture Collection which incorporate derivatives of the  $\lambda$  P<sub>L</sub> promoter include plasmid pHUB2, resident in *E. coli* strain JMB9 (ATCC 37092) and pPLc28, resident in *E. coli* RR1 (ATCC 53082).

Recombinant type II IL-1R proteins may also be expressed in yeast hosts, preferably from the *Saccharomyces* species, such as *S. cerevisiae*. Yeast of other genera, such as *Pichia* or *Kluyveromyces* may also be employed. Yeast vectors will generally contain an origin of replication from the 2 $\mu$  yeast plasmid or an autonomously replicating sequence (ARS), promoter, DNA encoding type II IL-1R, sequences for polyadenylation and transcription termination and a selection gene. Preferably, yeast vectors will include an origin of replication and selectable marker permitting transformation of both yeast and *E. coli*, e.g., the ampicillin resistance gene of *E. coli* and *S. cerevisiae* TRP1 or URA3 gene, which provides a selection marker for a mutant strain of yeast lacking the ability to grow in tryptophan, and a promoter derived from a highly expressed yeast gene to induce transcription of a structural sequence downstream. The presence of the TRP1 or URA3 lesion in the yeast host cell genome then provides an effective environment for detecting transformation by growth in the absence of tryptophan or uracil.

Suitable promoter sequences in yeast vectors include the promoters for metallothionein, 3-phosphoglycerate kinase (Hitzeman et al., *J. Biol. Chem.* 255:2073, 1980) or other glycolytic enzymes (Hess et al., *J. Adv. Enzyme Reg.* 7:149, 1968; and Holland et al., *Biochem.* 17:4900, 1978), such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase. Suitable vectors and promoters for use in yeast expression are further described in R. Hitzeman et al., EPA 73,657.

Preferred yeast vectors can be assembled using DNA sequences from pUC18 for selection and replication in *E. coli* (Amp<sup>r</sup> gene and origin of replication) and yeast DNA sequences including a glucose-repressible ADH2 promoter and  $\alpha$ -factor secretion leader. The ADH2 promoter has been described by Russell et al. (*J. Biol. Chem.* 258:2674, 1982) and Beier et al. (*Nature* 300:724, 1982). The yeast  $\alpha$ -factor leader, which directs secretion of heterologous proteins, can be inserted between the promoter and the structural gene to be expressed. See, e.g., Kurjan et al., *Cell* 30:933, 1982; and Bitter et al., *Proc. Natl. Acad. Sci. USA* 81:5330, 1984. The leader sequence may be modified to contain, near its 3' end, one or more useful restriction sites to facilitate fusion of the leader sequence to foreign genes.

Suitable yeast transformation protocols are known to those of skill in the art; an exemplary technique is described by Hinnen et al., *Proc. Natl. Acad. Sci. USA* 75:1929,

1978, selecting for Trp<sup>+</sup> transformants in a selective medium consisting of 0.67% yeast nitrogen base, 0.5% casamino acids, 2% glucose, 10 µg/ml adenine and 20 µg/ml uracil or URA<sup>+</sup> transformants in medium consisting of 0.67% YNB, with amino acids and bases as described by Sherman et al., *Laboratory Course Manual for Methods in Yeast Genetics*, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, 1986.

Host strains transformed by vectors comprising the ADH2 promoter may be grown for expression in a rich medium consisting of 1% yeast extract, 2% peptone, and 1% or 4% glucose supplemented with 80 µg/ml adenine and 80 µg/ml uracil. Derepression of the ADH2 promoter occurs upon exhaustion of medium glucose. Crude yeast supernatants are harvested by filtration and held at 4°C prior to further purification.

Various mammalian or insect cell culture systems are also advantageously employed to express recombinant protein. Expression of recombinant proteins in mammalian cells is particularly preferred because such proteins are generally correctly folded, appropriately modified and completely functional. Examples of suitable mammalian host cell lines include the COS-7 lines of monkey kidney cells, described by Gluzman (*Cell* 23:175, 1981), and other cell lines capable of expressing an appropriate vector including, for example, L cells, C127, 3T3, Chinese hamster ovary (CHO), HeLa and BHK cell lines. Mammalian expression vectors may comprise nontranscribed elements such as an origin of replication, a suitable promoter and enhancer linked to the gene to be expressed, and other 5' or 3' flanking nontranscribed sequences, and 5' or 3' nontranslated sequences, such as necessary ribosome binding sites, a polyadenylation site, splice donor and acceptor sites, and transcriptional termination sequences. Baculovirus systems for production of heterologous proteins in insect cells are reviewed by Luckow and Summers, *BioTechnology* 6:47 (1988).

The transcriptional and translational control sequences in expression vectors to be used in transforming vertebrate cells may be provided by viral sources. For example, commonly used promoters and enhancers are derived from Polyoma, Adenovirus 2, Simian Virus 40 (SV40), and human cytomegalovirus. DNA sequences derived from the SV40 viral genome, for example, SV40 origin, early and late promoter, enhancer, splice, and polyadenylation sites may be used to provide the other genetic elements required for expression of a heterologous DNA sequence. The early and late promoters are particularly useful because both are obtained easily from the virus as a fragment which also contains the SV40 viral origin of replication (Fiers et al., *Nature* 273:113, 1978). Smaller or larger SV40 fragments may also be used, provided the approximately 250 bp sequence extending from the *Hind* 3 site toward the *Bgl*1 site located in the viral origin of replication is included. Further, mammalian genomic type II IL-1R promoter, control and/or signal sequences may be utilized, provided such control sequences are compatible with the host

cell chosen. Additional details regarding the use of a mammalian high expression vector to produce a recombinant mammalian type II IL-1R are provided in Examples 2 below. Exemplary vectors can be constructed as disclosed by Okayama and Berg (*Mol. Cell. Biol.* 3:280, 1983).

A useful system for stable high level expression of mammalian receptor cDNAs in C127 murine mammary epithelial cells can be constructed substantially as described by Cosman et al. (*Mol. Immunol.* 23:935, 1986).

In preferred aspects of the present invention, recombinant expression vectors comprising type II IL-1R cDNAs are stably integrated into a host cell's DNA. Elevated levels of expression product is achieved by selecting for cell lines having amplified numbers of vector DNA. Cell lines having amplified numbers of vector DNA are selected, for example, by transforming a host cell with a vector comprising a DNA sequence which encodes an enzyme which is inhibited by a known drug. The vector may also comprise a DNA sequence which encodes a desired protein. Alternatively, the host cell may be co-transformed with a second vector which comprises the DNA sequence which encodes the desired protein. The transformed or co-transformed host cells are then cultured in increasing concentrations of the known drug, thereby selecting for drug-resistant cells. Such drug-resistant cells survive in increased concentrations of the toxic drug by overproduction of the enzyme which is inhibited by the drug, frequently as a result of amplification of the gene encoding the enzyme. Where drug resistance is caused by an increase in the copy number of the vector DNA encoding the inhibitable enzyme, there is a concomitant co-amplification of the vector DNA encoding the desired protein (e.g., type II IL-1R) in the host cell's DNA.

A preferred system for such co-amplification uses the gene for dihydrofolate reductase (DHFR), which can be inhibited by the drug methotrexate (MTX). To achieve co-amplification, a host cell which lacks an active gene encoding DHFR is either transformed with a vector which comprises DNA sequence encoding DHFR and a desired protein, or is co-transformed with a vector comprising a DNA sequence encoding DHFR and a vector comprising a DNA sequence encoding the desired protein. The transformed or co-transformed host cells are cultured in media containing increasing levels of MTX, and those cell lines which survive are selected.

A particularly preferred co-amplification system uses the gene for glutamine synthetase (GS), which is responsible for the synthesis of glutamine from glutamate and ammonia using the hydrolysis of ATP to ADP and phosphate to drive the reaction. GS is subject to inhibition by a variety of inhibitors, for example methionine sulfoximine (MSX). Thus, type II IL-1R can be expressed in high concentrations by co-amplifying cells transformed with a vector comprising the DNA sequence for GS and a desired

protein, or co-transformed with a vector comprising a DNA sequence encoding GS and a vector comprising a DNA sequence encoding the desired protein, culturing the host cells in media containing increasing levels of MSX and selecting for surviving cells. The GS co-amplification system, appropriate recombinant expression vectors and cells lines, are described in the following PCT applications: WO 87/04462, WO 89/01036, WO 89/10404 and WO 86/05807.

Recombinant proteins are preferably expressed by co-amplification of DHFR or GS in a mammalian host cell, such as Chinese Hamster Ovary (CHO) cells, or alternatively in a murine myeloma cell line, such as SP2/0-Ag14 or NS0 or a rat myeloma cell line, such as YB2/3.0-Ag20, disclosed in PCT applications WO/89/10404 and WO 86/05807.

A preferred eukaryotic vector for expression of type II IL-1R DNA is disclosed below in Example 2. This vector, referred to as pDC406, was derived from the mammalian high expression vector pDC201 and contains regulatory sequences from SV40, HIV and EBV.

#### Purification of Recombinant Type II IL-1R

Purified mammalian type II IL-1Rs or analogs are prepared by culturing suitable host/vector systems to express the recombinant translation products of the DNAs of the present invention, which are then purified from culture media or cell extracts.

For example, supernatants from systems which secrete recombinant soluble type II IL-1R protein into culture media can be first concentrated using a commercially available protein concentration filter, for example, an Amicon or Millipore Pellicon ultrafiltration unit. Following the concentration step, the concentrate can be applied to a suitable purification matrix. For example, a suitable affinity matrix can comprise an IL-1 or lectin or antibody molecule bound to a suitable support. Alternatively, an anion exchange resin can be employed, for example, a matrix or substrate having pendant diethylaminoethyl (DEAE) groups. The matrices can be acrylamide, agarose, dextran, cellulose or other types commonly employed in protein purification. Alternatively, a cation exchange step can be employed. Suitable cation exchangers include various insoluble matrices comprising sulfopropyl or carboxymethyl groups. Sulfopropyl groups are preferred.

Finally, one or more reversed-phase high performance liquid chromatography (RP-HPLC) steps employing hydrophobic RP-HPLC media, e.g., silica gel having pendant methyl or other aliphatic groups, can be employed to further purify a type II IL-1R composition. Some or all of the foregoing purification steps, in various combinations, can also be employed to provide a homogeneous recombinant protein.

Recombinant protein produced in bacterial culture is usually isolated by initial extraction from cell pellets, followed by one or more concentration, salting-out, aqueous

ion exchange or size exclusion chromatography steps. Finally, high performance liquid chromatography (HPLC) can be employed for final purification steps. Microbial cells employed in expression of recombinant mammalian type II IL-1R can be disrupted by any convenient method, including freeze-thaw cycling, sonication, mechanical disruption, or use of cell lysing agents.

Fermentation of yeast which express soluble mammalian type II IL-1R as a secreted protein greatly simplifies purification. Secreted recombinant protein resulting from a large-scale fermentation can be purified by methods analogous to those disclosed by Urdal et al. (*J. Chromatog.* 296:171, 1984). This reference describes two sequential, reversed-phase HPLC steps for purification of recombinant human GM-CSF on a preparative HPLC column.

Human type II IL-1R synthesized in recombinant culture is characterized by the presence of non-human cell components, including proteins, in amounts and of a character which depend upon the purification steps taken to recover human type II IL-1R from the culture. These components ordinarily will be of yeast, prokaryotic or non-human higher eukaryotic origin and preferably are present in innocuous contaminant quantities, on the order of less than about 1 percent by weight. Further, recombinant cell culture enables the production of type II IL-1R free of proteins which may be normally associated with type II IL-1R as it is found in nature in its species of origin, e.g. in cells, cell exudates or body fluids.

#### Therapeutic Administration of Recombinant Soluble Type II IL-1R

The present invention provides methods of using therapeutic compositions comprising an effective amount of soluble type II IL-1R proteins and a suitable diluent and carrier, and methods for suppressing IL-1-dependent immune responses in humans comprising administering an effective amount of soluble type II IL-1R protein.

For therapeutic use, purified soluble type II IL-1R protein is administered to a patient, preferably a human, for treatment in a manner appropriate to the indication. Thus, for example, soluble type II IL-1R protein compositions can be administered by bolus injection, continuous infusion, sustained release from implants, or other suitable technique. Typically, a soluble type II IL-1R therapeutic agent will be administered in the form of a composition comprising purified protein in conjunction with physiologically acceptable carriers, excipients or diluents. Such carriers will be nontoxic to recipients at the dosages and concentrations employed. Ordinarily, the preparation of such compositions entails combining the type II IL-1R with buffers, antioxidants such as ascorbic acid, low molecular weight (less than about 10 residues) polypeptides, proteins, amino acids, carbohydrates including glucose, sucrose or dextrans, chelating agents such as EDTA,

glutathione and other stabilizers and excipients. Neutral buffered saline or saline mixed with conspecific serum albumin are exemplary appropriate diluents. Preferably, product is formulated as a lyophilizate using appropriate excipient solutions (e.g., sucrose) as diluents. Appropriate dosages can be determined in trials; generally, shuIL-1R dosages of from about 1 ng/kg/day to about 10 mg/kg/day, and more preferably from about 500 µg/kg/day to about 5 mg/kg/day, are expected to induce a biological effect.

Because IL-1R-I and type II IL-1R proteins both bind to IL-1, soluble type II IL-1R proteins are expected to have similar, if not identical, therapeutic activities. For example, soluble human type II IL-1R can be administered, for example, for the purpose of suppressing immune responses in a human. A variety of diseases or conditions are caused by an immune response to alloantigen, including allograft rejection and graft-versus-host reaction. In alloantigen-induced immune responses, shuIL-1R suppresses lymphoproliferation and inflammation which result upon activation of T cells. shuIL-1R can therefore be used to effectively suppress alloantigen-induced immune responses in the clinical treatment of, for example, rejection of allografts (such as skin, kidney, and heart transplants), and graft-versus-host reactions in patients who have received bone marrow transplants.

Soluble human type II IL-1R can also be used in clinical treatment of autoimmune dysfunctions, such as rheumatoid arthritis, diabetes and multiple sclerosis, which are dependent upon the activation of T cells against antigens not recognized as being indigenous to the host.

The following examples are offered by way of illustration, and not by way of limitation.

## **EXAMPLES**

### **Example 1**

#### **Isolation of cDNA Encoding Human Type II IL-1R by Direct Expression of Active Protein in CV-1/EBNA-1 Cells**

A. Radiolabeling of rIL-1β. Recombinant human IL-1β was prepared by expression in *E. coli* and purification to homogeneity as described by Kronheim et al. (*BioTechnology* 4:1078, 1986). The IL-1β was labeled with di-iodo (<sup>125</sup>I) Bolton-Hunter reagent (New England Nuclear, Glenolden, PA). Ten micrograms (0.57 nmol) of protein in 10 µL of phosphate (0.015 mol/L)-buffered saline (PBS; 0.15 mol/L), pH 7.2, was mixed with 10 µL of sodium borate (0.1 mol/L)-buffered saline (0.15 mol/L), pH 8.5, and reacted with 1 mCi (0.23 nmol) of Bolton-Hunter reagent according to the manufacturer's instructions for 12 hours at 8°C. Subsequently, 30 µL of 2% gelatin and 5 µL of 1 mol/L glycine ethyl ester were added, and the protein was separated from unreacted Bolton-

Hunter reagent on a 1 mL bed volume Biogel™ P6 column (Bio-Rad Laboratories, Richmond, CA). Routinely, 50% to 60% incorporation of label was observed. Radiiodination yielded specific activities in the range of  $1 \times 10^{15}$  to  $5 \times 10^{15}$  cpm/mmol-1 (0.4 to 2 atoms I per molecule protein), and sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) revealed a single labeled polypeptide of 17.5 kD, consistent with previously reported values for IL-1. The labeled protein was greater than 98% TCA precipitable, indicating that the  $^{125}\text{I}$  was covalently bound to protein.

**B. Construction and Screening of CB23 cDNA library.** A CB23 library was constructed and screened by direct expression of pooled cDNA clones in the monkey kidney cell line CV-1/EBNA-1 (which was derived by transfection of the CV-1 cell line with the gene encoding EBNA-1, as described below) using a mammalian expression vector (pDC406) that includes regulatory sequences from SV40, human immunodeficiency virus (HIV), and Epstein-Barr virus (EBV). The CV-1/EBNA-1 cell line constitutively expresses EBV nuclear antigen-1 driven from the human cytomegalovirus (CMV) immediate-early enhancer/promoter and therefore allows the episomal replication of expression vectors such as pDC406 that contain the EBV origin of replication. The expression vector used was pDC406, a derivative of HAV-EO, described by Dower et al., *J. Immunol.* 142:4314, 1989), which is in turn a derivative of pDC201 and allows high level expression in the CV-1/EBNA-1 cell line. pDC406 differs from HAV-EO (Dower et al., *supra*) by the deletion of the intron present in the adenovirus 2 tripartite leader sequence in HAV-EO (see description of pDC303 below).

The CB23 cDNA library was constructed by reverse transcription of poly(A)<sup>+</sup> mRNA isolated from total RNA extracted from the human B cell lymphoblastoid line CB23 (Benjamin & Dower, *Blood* 75:2017, 1990) substantially as described by Ausubel et al., eds., *Current Protocols in Molecular Biology*, Vol. 1, 1987. The CB23 cell line is an EBV-transformed cord blood (CB) lymphocyte cell line, which was derived by using the methods described by Benjamin et al., *Proc. Natl. Acad. Sci. USA* 81:3547, 1984. Poly(A)<sup>+</sup> mRNA was isolated by oligo dT cellulose chromatography and double-stranded cDNA was made substantially as described by Gubler and Hoffman, *Gene* 25:263, 1983. Briefly, the poly(A)<sup>+</sup> mRNA was converted to an RNA-cDNA hybrid with reverse transcriptase using random hexanucleotides as a primer. The RNA-cDNA hybrid was then converted into double-stranded cDNA using RNAase H in combination with DNA polymerase I. The resulting double stranded cDNA was blunt-ended with T4 DNA polymerase. The following two unkinased oligonucleotides were annealed and blunt end ligated with DNA ligase to the ends of the resulting blunt-ended cDNA as described by Haymerle, et al., *Nucleic Acids Research*, 14: 8615, 1986.

```

SEQ ID NO:3      5'- TCG ACT GGA ACG AGA CGA CCT GCT -3'
SEQ ID NO:4      3'-      GA CCT TGC TCT GCT GGA CGA -5'
                  <SalI>

```

In this case only the 24-mer oligo will ligate onto the cDNA. The non-ligated oligos were removed by gel filtration chromatography at 68°C, leaving 24 nucleotide non-self-complementary overhangs on the cDNA. The same procedure was used to convert the 5' ends of *SalI*-cut mammalian expression vector pDC406 to 24 nucleotide overhangs complementary to those added to the cDNA. Optimal proportions of adapted vector and cDNA were ligated in the presence of T4 polynucleotide kinase. Dialyzed ligation mixtures were electroporated into *E. coli* strain DH5α. Approximately  $3.9 \times 10^6$  clones were generated and plated in pools of approximately 3,000. A sample of each pool was used to prepare frozen glycerol stocks and a sample was used to obtain a pool of plasmid DNA.

The pooled DNA was then used to transfect a sub-confluent layer of monkey CV-1/EBNA-1 cells using DEAE-dextran followed by chloroquine treatment, similar to that described by Luthman et al., *Nucl. Acids Res.* 11:1295 (1983) and McCutchan et al., *J. Natl. Cancer Inst.* 41:351 (1986). CV-1/EBNA-1 cells were derived as follows. The CV-1/EBNA-1 cell line constitutively expresses EBV nuclear antigen-1 driven from the CMV immediate-early enhancer/promoter. The African Green Monkey kidney cell line, CV-1 (ATCC CCL 70, was cotransfected with 5 µg of pSV2gpt (Mulligan & Berg, *Proc. Natl. Acad. Sci. USA* 78:2072, 1981) and 25 µg of pDC303/EBNA-1 using a calcium phosphate coprecipitation technique (Ausubel et al., eds., *Current Protocols in Molecular Biology*, Wiley, New York, 1987). pDC303/EBNA-1 was constructed from pDC302 (Mosley et al., *Cell* 59:335, 1989) in two steps. First, the intron present in the adenovirus tripartite leader sequence was deleted by replacing a PvuII to ScaI fragment spanning the intron with the following synthetic oligonucleotide pair to create plasmid pDC303:

SEQ ID NO:5 5'-CTGTTGGGCTCGCGGTTGAGGACAACTCTTCGCGGTCTTTCCAGT-3'

SEQ ID NO:6 3'-GACAACCCGAGCGCCAACTCCTGTTTGAGAAGCGCCAGAAAGGTCA-5'

Second, a HindIII-AhaII restriction fragment encoding Epstein-Barr virus nuclear antigen I (EBNA-1), and consisting essentially of EBV coordinates 107,932 to 109,894 (Baer et al., *Nature* 310:207, 1984), was then inserted into the multiple cloning site of pDC303 to create the plasmid pDC303/EBNA-1. The transfected cells were grown in the presence of hypoxanthine, aminopterin, thymidine, xanthine, and mycophenolic acid according to standard methods (Ausubel et al., *supra*; Mulligan & Berg, *supra*) to select for the cells that had stably incorporated the transfected plasmids. The resulting drug resistant colonies were isolated and expanded individually into cell lines for analysis. The cell lines were

screened for the expression of functional EBNA-1. One cell line, clone 68, was found to express EBNA-1 using this assay, and was designated CV-1/EBNA-1.

In order to transfect the CV-1/EBNA-1 cells with the cDNA library, the cells were maintained in complete medium (Dulbecco's modified Eagle's media (DMEM) containing 10% (v/v) fetal calf serum (FCS), 50 U/ml penicillin, 50 U/ml streptomycin, 2 mM L-glutamine) and were plated at a density of  $2 \times 10^5$  cells/well in either 6 well dishes (Falcon) or single well chambered slides (Lab-Tek). Both dishes and slides were pretreated with 1 ml human fibronectin (10 ug/ml in PBS) for 30 minutes followed by 1 wash with PBS. Media was removed from the adherent cell layer and replaced with 1.5 ml complete medium containing 66.6  $\mu$ M chloroquine sulfate. 0.2 mls of DNA solution (2  $\mu$ g DNA, 0.5 mg/ml DEAE-dextran in complete medium containing chloroquine) was then added to the cells and incubated for 5 hours. Following the incubation, the media was removed and the cells shocked by addition of complete medium containing 10% DMSO for 2.5 to 20 minutes followed by replacement of the solution with fresh complete medium. The cells were grown in culture to permit transient expression of the inserted sequences. These conditions led to an 80% transfection frequency in surviving CV-1/EBNA-1 cells.

After 48 to 72 hours, transfected monolayers of CV-1/EBNA cells were assayed for expression of IL-1 binding proteins by binding radioiodinated IL-1 $\beta$  prepared as described above by slide autoradiography. Transfected CV-1/EBNA-1 cells were washed once with binding medium (RPMI medium 1640 containing 25 mg/ml bovine serum albumin (BSA), 2 mg/ml sodium azide, 20 mM HEPES, pH 7.2, and 50 mg/ml nonfat dry milk (NFDM)) and incubated for 2 hours at 4°C with 1 ml binding medium + NFDM containing  $3 \times 10^{-9}$  M  $^{125}$ I-IL-1 $\beta$ . After incubation, cells in the chambered slides were washed three times with binding buffer + NFDM, followed by 2 washes with PBS, pH 7.3, to remove unbound  $^{125}$ I-IL-1 $\beta$ . The cells were fixed by incubating for 30 minutes at room temperature in 10% glutaraldehyde in PBS, pH 7.3, washed twice in PBS, and air dried. The slides were dipped in Kodak GTNB-2 photographic emulsion (6x dilution in water) and exposed in the dark for 48 hours to 7 days at 4°C in a light proof box. The slides were then developed for approximately 5 minutes in Kodak D19 developer (40 g/500 ml water), rinsed in water and fixed in Agfa G433C fixer. The slides were individually examined with a microscope at 25-40x magnification and positive cells expressing type II IL-1R were identified by the presence of autoradiographic silver grains against a light background.

Cells in the 6 well plates were washed once with binding buffer + NFDM followed by 3 washings with PBS, pH 7.3, to remove unbound  $^{125}$ I-IL-1 $\beta$ . The bound cells were then trypsinized to remove them from the plate and bound  $^{125}$ I-IL-1 $\beta$  were counted on a beta counter.

### Example 2

The 5' primer corresponds to nucleotides 31-51 from the untranslated region of human type II IL-1R clone 75 (SEQ ID NO:1) with a 5' add-on of a SalI restriction site; this nucleotide sequence is capable of annealing to the (-) strand complementary to nucleotides 31-51 of human clone 75. The 3' primer is complementary to nucleotides 1191-1172 (which includes anti-sense nucleotides encoding 3 amino acids of human type II IL-1R clone 75 (SEQ ID NO:1) and has a 5' add-on of a NotI restriction site and a stop codon.

The following PCR reagents were added to a 1.5 ml Eppendorf microfuge tube: 10  $\mu$ l of 10X PCR buffer (500 mM KCl, 100 mM Tris-HCl, pH 8.3 at 25°C, 15 mM MgCl<sub>2</sub>, and 1 mg/ml gelatin) (Perkin-Elmer Cetus, Norwalk, CN), 10  $\mu$ l of a 2mM solution containing each dNTP (2 mM dATP, 2 mM dCTP, 2 mM dGTP and 2 mM dTTP), 2.5 units (0.5  $\mu$ l of standard 5000 units/ml solution) of *Taq* DNA polymerase (Perkin-Elmer Cetus), 50 ng of template DNA and 5  $\mu$ l of a 20  $\mu$ M solution of each of the above oligonucleotide primers and 74.5  $\mu$ l water to a final volume of 100  $\mu$ l. The final mixture was then overlaid with 100  $\mu$ l paraffin oil. PCR was carried out using a DNA thermal cycler (Ericomp, San Diego, CA) by initially denaturing the template at 94° for 90 seconds, reannealing at 55° for 75 seconds and extending the cDNA at 72° for 150 seconds. PCR was carried out for an additional 20 cycles of amplification using a step program (denaturation at 94°, 25 sec; annealing at 55°, 45 sec; extension at 72°, 150 sec.), followed by a 5 minute extension at 72°.

The sample was removed from the paraffin oil and DNA extracted by phenol-chloroform extraction and spun column chromatography over G-50 (Boehringer Mannheim). A 10  $\mu$ l aliquot of the extracted DNA was separated by electrophoresis on 1% SeaKem™ agarose (FMC BioProducts, Rockland, ME) and stained with ethidium bromide to confirm that the DNA fragment size was consistent with the predicted product.

20  $\mu$ l of the PCR-amplified cDNA products were then digested with *Sal*I and *Not*I restriction enzymes using standard procedures. The *Sal*I/*Not*I restriction fragment was then separated on a 1.2% Seaplaque™ low gelling temperature (LGT) agarose, and the band representing the fragment was isolated. The fragment was ligated into the pDC406 vector by a standard "in gel" ligation method, and the vector was transfected into CV1-EBNA cells and expressed as described above in Example 1.

### Example 3

#### Isolation of cDNAs Encoding Murine Type II IL-1R

Murine type II IL-1R cDNAs were isolated from a cDNA library made from the murine pre-B cell line 70Z/3 (ATCC TIB 158), by cross species hybridization with a human Type II IL-1R probe. A cDNA library was constructed in a  $\lambda$  phage vector using  $\lambda$ gt10 arms and packaged in vitro (Gigapack®, Stratagene, San Diego) according to the manufacturer's instructions. A double-stranded human Type II IL-1R probe was produced by excising an approximately 1.35 kb *Sal*I restriction fragment of the human type II IL-1R clone 75 and <sup>32</sup>P-labelling the cDNA using random primers (Boehringer-Mannheim). The murine cDNA library was amplified once and a total of 5x10<sup>5</sup> plaques were screened with the human probe in 35% formamide (5xSSC, 42°C). Several murine type II IL-1R cDNA clones (including clone  $\lambda$ 2) were isolated; however, none of the clones appeared to be full-

length. Nucleotide sequence information obtained from the partial clones was used to clone a full-length murine type II IL-1R cDNA as follows.

A full-length cDNA clone encoding murine type II IL-1R was isolated by the method of Rapid Amplification of cDNA Ends (RACE) described by Frohman et al., *Proc. Natl. Acad. Sci. USA* 85:8998, 1988, using RNA from the murine pre-B cell line 70Z/3. Briefly, the RACE method uses PCR to amplify copies of a region of cDNA between a known point in the cDNA transcript (determined from nucleotide sequence obtained as described above) and the 3' end. An adaptor-primer having a sequence containing 17 dT base pairs and an adaptor sequence containing three endonuclease recognition sites (to place convenient restriction sites at the 3' end of the cDNA) is used to reverse transcribe a population of mRNA and produce (-) strand cDNA. A primer complementary to a known stretch of sequence in the 5' untranslated region of the murine type II IL-1R clone 2 cDNA, described above, and oriented in the 3' direction is annealed with the (-) strand cDNA and extended to generate a complementary (+) strand cDNA. The resulting double-strand cDNA is amplified by PCR using primers that anneal to the natural 5'-end and synthetic 3'-end poly(A) tail. Details of the RACE procedure are as follows.

The following PCR oligonucleotide primers (d(T)<sub>17</sub> adaptor-primer, 5' amplification primer and 3' amplification primer, respectively) were first constructed:

SEQ ID NO:9	5'-CTGCAGGCGGCCGCGGATCC(T) <sub>17</sub> -3' <PstI><-NotI-><BamHI>
SEQ ID NO:10	5'-GCGTCGACGGCAAGAAGCAGCAAGGTAC-3' <SalI>\15 \34
SEQ ID NO:11	5'-CTGCAGGCGGCCGCGGATCC-3' <PstI><-NotI-><BamHI>

Briefly, the d(T)<sub>17</sub> adapter-primer (SEQ ID NO:9) contains nucleotide sequence anneals to the poly(A)<sup>+</sup> region of a population mRNA transcripts and is used to generated (-) strand cDNA reverse transcripts from mRNA; it also contains endonuclease restriction sites for PstI, NotI and BamHI to be introduced into the DNA being amplified by PCR. The 5' amplification primer (SEQ ID NO:10) corresponds to nucleotides 15-34 from the 5' untranslated region of murine type II IL-1R clone λ2 with a 5' add-on of a SalI restriction site; this nucleotide sequence anneals to the (-) strand cDNA generated by reverse transcription with the d(T)<sub>17</sub> adaptor-primer and is extended to generate (+) strand cDNA. The 3' primer (SEQ ID NO:11) anneals to the (+) strand DNA having the above

endonuclease restriction sites and is extended to generate a double-stranded full-length cDNA encoding murine type II IL-1R, which can then be amplified by a standard PCR reaction. Details of the PCR procedure are as follows.

Poly(A)<sup>+</sup> mRNA was isolated by oligo dT cellulose chromatography from total RNA extracted from 70Z/3 cells using standard methods described by Maniatis et al., *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor Lab., Cold Spring Harbor, NY, 1982) and reverse transcribed as follows. Approximately 1 µg of poly(A)<sup>+</sup> mRNA in 16.5 µl of water was heated at 68°C for 3 minutes and then quenched on ice, and added to 2 µl of 10X RTC buffer (500 mM Tris-HCl, pH 8.7 at 22°C, 60 mM MgCl<sub>2</sub>, 400 mM KCl, 10 mM DTT, each dNTP at 10 mM), 10 units of RNasin (Promega Biotech), 0.5 µg of d(T)<sub>17</sub>-adapter primer and 10 units of AMV reverse transcriptase (Life Sciences) in a total volume of 20 µl, and incubated for a period of 2 hours at 42°C to reverse transcribe the mRNA and synthesize a pool of cDNA. The reaction mixture was diluted to 1 ml with TE buffer (10 mM Tris-HCl, pH 7.5, 1 mM EDTA) and stored at 4°C overnight.

Approximately 1 or 5 µl of the cDNA pool was combined with 5 µl of a 20 µM solution of the 5' amplification primer, containing sequence corresponding to the sequence of nucleotides 15-34 of murine type II IL-1R clone λ2, 5 µl of a 20 µM solution of the 3' amplification primer, 10 µl of 10X PCR buffer (500 mM KCl, 100 mM Tris-HCl (pH 8.4, 20°C), 14 mM MgCl<sub>2</sub>, and 1 mg/ml gelatin), 4 µl of 5mM each dNTP (containing 5 mM dATP, 5mM dCTP, 5mM dGTP and 5mM dTTP), 2.5 units (0.5 µl of standard 5000 units/ml solution) of *Taq* DNA polymerase (Perkin-Elmer Cetus Instruments), diluted to a volume of 100 µl. The final mixture was then overlaid with 100 µl paraffin oil. PCR was carried out using a DNA thermal cycler (Perkin-Elmer/Cetus) by initially denaturing the template at 94° for 90 seconds, reannealing at 64° for 75 seconds and extending the cDNA at 72° for 150 seconds. PCR was carried out for an additional 25 cycles of amplification using the following step program (denaturation at 94° for 25 sec; annealing at 55° for 45 sec; extension at 72° for 150 sec.), followed by a 7 minute final extension at 72°.

The sample was removed from the paraffin oil and DNA extracted by phenol-chloroform extraction and spun column chromatography over G-50 (Boehringer Mannheim). A 10 µl aliquot of the extracted DNA was separated by electrophoresis on 1% SeaKem™ agarose (FMC BioProducts, Rockland, ME) and stained with ethidium bromide to confirm that the DNA fragment size was consistent with the predicted product. The gel was then blotted and probed with a 5' 610bp EcoRI fragment of murine type II IL-1R clone λ2 from above to confirm that the band contained DNA encoding murine type II IL-1R.

The PCR-amplified cDNA products were then concentrated by centrifugation in an Eppendorf microfuge at full speed for 20 min., followed by ethanol precipitation in 1/10 volume sodium acetate (3 M) and 2.5 volume ethanol. 30 µl of the concentrate was

digested with Sall and NotI restriction enzymes using standard procedures. The Sall/NotI restriction fragment was then separated on a 1.2% LGT agarose gel, and the band representing the fragment was isolated. The restriction fragments were then purified from the agarose using GeneClean™ (Bio-101, La Jolla, CA).

The resulting purified restriction fragment was ligated into the pDC406 vector, which was then transfected into CV1-EBNA cells and expressed as described above in Example 1.

The Sequence Listing setting forth the nucleotide (SEQ ID No:12) and predicted amino acid sequences (SEQ ID No:12 and SEQ ID NO:13) and associated information appears at the end of the specification immediately prior to the claims.

#### Example 4

##### Construction and Expression of cDNAs Encoding Murine Soluble Type II IL-1R

A cDNA encoding soluble murine type II IL-1R (having the sequence of amino acids -13-345 of SEQ ID NO:12) was constructed by PCR amplification 70Z/3 poly(A)<sup>+</sup> mRNA as a template and the following procedure as described for the full length clone encoding murine type II IL-1R. The following PCR oligonucleotide primers (d(T)<sub>17</sub> adaptor-primer, 5' amplification primer and 3' amplification primer, respectively) were constructed:

SEQ ID NO:9	5'-CTGCAGGCGGCCGCGGATCC(T) <sub>17</sub> -3' <PstI><-NotI--><BamHI>
SEQ ID NO:10	5'-GCGTCGACGGCAAGAAGCAGCAAGGTAC-3' <SalI>\15 \34
SEQ ID NO:14	5'-GCGCGGCCGCGCTAGGAAGAGACTTCTTTGACTGTGG-3' <--NotI-->EndSerSerValGluLysValThrThr

The d(T)<sub>17</sub> adaptor-primer and 5' amplification primer are identical with SEQ ID NO:9 and SEQ ID NO:10, described in Example 5. The 3' end of SEQ ID NO:12 is complementary to nucleotides 1145-1166 of SEQ ID NO:12 and has a 5' add-on of a NotI restriction site and a stop codon.

A pool of cDNA was synthesized from poly(A)<sup>+</sup> mRNA using the d(T)<sub>17</sub> adaptor-primer as described in Example 3. To a 1.5 ml Eppendorf microfuge tube was added approximately 1 µl of the cDNA pool, 5 µl of a 20 µM solution of the 5' amplification primer, 5 µl of a 20 µM solution of the 3' amplification primer, 10 µL of 10X PCR buffer

(500 mM KCl, 100 mM Tris-HCl (pH 8.4 at 20°C), 14 mM MgCl<sub>2</sub>, and 1 mg/ml gelatin), 4 µl of 5mM each of dNTP (containing 5 mM dATP, 5mM dCTP, 5mM dGTP and 5mM dTTP), 2.5 units (0.5 µl of standard 5000 units/ml solution) of *Taq* DNA polymerase (Perkin-Elmer Cetus Instruments), diluted with 75.4 µl water to a volume of 100 µl. The final mixture was then overlaid with 100 µl paraffin oil. PCR was carried out using a DNA thermal cycler (Ericomp) by initially denaturing the template at 94° for 90 seconds, reannealing at 55° for 75 seconds and extending the cDNA at 72° for 150 seconds. PCR was carried out for an additional 20 cycles of amplification using the following step program (denaturation at 94° for 25 sec; annealing at 55° for 45 sec; extension at 72° for 150 sec.), followed by a 7 minute final extension at 72°.

The sample was removed from the paraffin oil and DNA extracted by phenol-chloroform extraction and spun column chromatography over G-50 (Boehringer Mannheim). A 10 µl aliquot of the extracted DNA was separated by electrophoresis on 1% SeaKem™ agarose (FMC BioProducts, Rockland, ME) and stained with ethidium bromide to confirm that the DNA fragment size was consistent with the predicted product.

The PCR-amplified cDNA products were then concentrated by centrifugation in an Eppendorf microfuge at full speed for 20 min., followed by ethanol precipitation in 1/10 volume sodium acetate (3 M) and 2.5 volume ethanol. 50 µl was digested with *Sall* and *NotI* restriction enzymes using standard procedures. The *Sall/NotI* restriction fragment was then separated on a 1.2% Seaplaque LGT agarose gel, and the band representing the fragment was isolated. The restriction fragment was then purified from the isolated band using the following freeze/thaw method. The band from the gel was split into two 175 µl fragments and placed into two 1.5 ml Eppendorf microfuge tubes. 500 µl of isolation buffer (0.15 M NaCl, 10 mM Tris, pH 8.0, 1mM EDTA) was added to each tube and the tubes heated to 68°C to melt the gel. The gels were then frozen on dry ice for 10 minutes, thawed at room temperature and centrifuged at 4°C for 30 minutes. Supernatants were then removed and placed in a new tube, suspended in 2 mL ethanol, and centrifuged at 4°C for an additional 30 minutes to form a DNA pellet. The DNA pellet was washed with 70% ethanol, centrifuged for 5 minutes, removed from the tube and resuspended in 20 µl TE buffer.

The resulting purified restriction fragments were then ligated into the pDC406 vector. A sample of the ligation was transformed into DH5α and colonies were analyzed to check for correct plasmids. The vector was then transfected into COS-7 cells and expressed as described above in Example 1.

### Example 5

#### Type II IL-1R Binding Studies

The binding inhibition constant of recombinant human type II IL-1R, expressed and purified as described in Example 1 above, was determined by inhibition binding assays in which varying concentrations of a competitor (IL-1 $\beta$  or IL-1 $\alpha$ ) was incubated with a constant amount of radiolabeled IL-1 $\beta$  or IL-1 $\alpha$  and cells expressing the type II IL-1R. The competitor binds to the receptor and prevents the radiolabeled ligand from binding to the receptor. Binding assays were performed by a phthalate oil separation method essentially as describe by Dower et al., *J. Immunol.* 132:751, 1984 and Park et al., *J. Biol. Chem.* 261:4177, 1986. Briefly, CV1/EBNA cells were incubated in six-well plates (Costar, Cambridge, MA) at 4°C for 2 hours with <sup>125</sup>I-IL-1 $\beta$  in 1 ml binding medium (Roswell Park Memorial Institute (RPMI) 1640 medium containing 2% BSA, 20 mM Hepes buffer, and 0.2% sodium azide, pH 7.2). Sodium azide was included to inhibit internalization and degradation of <sup>125</sup>I-IL-1 by cells at 37°C. The plates were incubated on a gyratory shaker for 1 hour at 37°C. Replicate aliquots of the incubation mixture were then transferred to polyethylene centrifuge tubes containing a phthalate oil mixture comprising 1.5 parts dibutylphthalate, to 1 part bis(s-ethylhexyl)phthalate. Control tubes containing a 100X molar excess of unlabeled IL-1 $\beta$  were also included to determine non-specific binding. The cells with bound <sup>125</sup>I-IL-1 were separated from unbound <sup>125</sup>I-IL-1 by centrifugation for 5 minutes at 15,000X g in an Eppendorf Microfuge. The radioactivity associated with the cells was then determined on a gamma counter. This assay (using unlabeled human IL-1 $\beta$  as a competitor to inhibit binding of <sup>125</sup>I-IL-1 $\beta$  to type II IL-1R) indicated that the full length human type II IL-1R exhibits biphasic binding to IL-1 $\beta$  with a K<sub>I1</sub> of approximately  $19 \pm 8 \times 10^9$  and K<sub>I2</sub> of approximately  $0.2 \pm 0.002 \times 10^9$ . Using unlabeled human IL-1 $\beta$  to inhibit binding of <sup>125</sup>I-IL-1 $\alpha$  to type II IL-1R, the full length human type II IL-1R exhibited biphasic binding to IL-1 $\beta$  with a K<sub>I1</sub> of approximately  $2.0 \pm 1 \times 10^9$  and K<sub>I2</sub> of approximately  $0.013 \pm 0.003 \times 10^9$ .

The binding inhibition constant of the soluble human type II IL-1R, expressed and purified as described in Example 2 above, is determined by a inhibition binding assay in which varying concentrations of an IL-1 $\beta$  competitor is incubated with a constant amount of radiolabeled I-IL-1 $\beta$  and CB23 cells (an Epstein Barr virus transformed cord blood B lymphocyte cell line) expressing the type II IL-1R. Binding assays were also performed by a phthalate oil separation method essentially as describe by Dower et al., *J. Immunol.* 132:751, 1984 and Park et al., *J. Biol. Chem.* 261:4177, 1986. Briefly, COS-7 cells were transfected with the expression vector pDC406 containing a cDNA encoding the soluble human type II IL-1R described above. Supernatants from the COS cells were harvested 3

days after transfection and serially diluted in binding medium (Roswell Park Memorial Institute (RPMI) 1640 medium containing 2% BSA, 20 mM Hepes buffer, and 0.2% sodium azide, pH 7.2) in 6 well plates to a volume of 50  $\mu$ l/well. The supernatants were incubated with 50  $\mu$ l of  $9 \times 10^{-10}$  M  $^{125}$ I-IL-1 $\beta$  plus  $2.5 \times 10^6$  CB23 cells at 8°C for 2 hours with agitation. Duplicate 60  $\mu$ l aliquots of the incubation mixture were then transferred to polyethylene centrifuge tubes containing a phthalate oil mixture comprising 1.5 parts dibutylphthalate, to 1 part bis(s-ethylhexyl)phthalate. A negative control tube containing  $3 \times 10^{-6}$  M unlabeled IL-1 $\beta$  was also included to determine non-specific binding (100% inhibition) and a positive control tube containing 50 ml binding medium with only radiolabeled IL-1 $\beta$  was included to determine maximum binding. The cells with bound  $^{125}$ I-IL-1 $\beta$  were separated from unbound  $^{125}$ I-IL-1 $\beta$  by centrifugation for 5 minutes at 15,000 X g in an Eppendorf Microfuge. Supernatants containing unbound  $^{125}$ I-IL-1 $\beta$  were discarded and the cells were carefully rinsed with ice-cold binding medium. The cells were then incubated in 1 ml of trypsin-EDTA at 37°C for 15 minutes and cells were harvested. The radioactivity of the cells was then determined on a gamma counter. This inhibition binding assay (using soluble human type II IL-1R to inhibit binding of IL-1 $\beta$ ) indicated that the soluble human type II IL-1R has a  $K_I$  of approximately  $3.5 \times 10^9$  M $^{-1}$ . Inhibition of IL-1 $\alpha$  binding by soluble human type II IL-1R using the same procedure indicated that soluble human type II IL-1R has a  $K_I$  of  $1.4 \times 10^8$  M $^{-1}$ .

Murine type II IL-1R exhibits biphasic binding to IL-1 $\beta$  with a  $K_{I1}$  of  $0.8 \times 10^9$  and a  $K_{I2}$  of less than  $0.01 \times 10^9$ .

### Example 6

#### Type II IL-1R Affinity Crosslinking Studies

Affinity crosslinking studies were performed essentially as described by Park et al., *Proc. Natl. Acad. Sci. USA* 84:1669, 1987. Recombinant human IL-1 $\alpha$  and IL-1 $\beta$  used in the assays were expressed, purified and labeled as described previously (Dower et al., *J. Exp. Med.* 162:501, 1985; Dower et al., *Nature* 324:266, 1986). Recombinant human IL-1 receptor antagonist (IL-1ra) was cloned using the cDNA sequence published by Eisenberg et al., *Nature* 343:341, 1990, expressed by transient transfection in COS cells, and purified by affinity chromatography on a column of soluble human type I IL-1R coupled to affigel, as described by Dower et al., *J. Immunol.* 143:4314, 1989, and eluted at low pH.

Briefly, CV1/EBNA cells ( $4 \times 10^7$ /ml) expressing recombinant type II IL-1R were incubated with  $^{125}$ I-IL-1 $\alpha$  or  $^{125}$ I-IL-1 $\beta$  (1 nM) at 4°C in the presence and absence of 1  $\mu$ M excess of unlabeled IL-1 as a specificity control for 2 hours. The cells were then washed and bis(sulfosuccinimidyl)suberate was added to a final concentration of 0.1 mg/ml. After

30 min. at 25°C, the cells were washed and resuspended in 100 µl of phosphate-buffered saline (PBS)/1% Triton containing 2 mM leupeptin, 2 mM o-phenanthroline, and 2 mM EGTA to prevent proteolysis. Aliquots of the extract supernatants containing equal amounts (CPM) of  $^{125}\text{I}$ -IL-1 and equal volumes of the specificity controls, were analyzed by SDS/PAGE on a 10% gel using standard techniques.

Figure 4 shows the results of affinity crosslinking studies conducted as described above, using radiolabeled IL-1 $\alpha$  and IL-1 $\beta$ , to compare the sizes of the recombinant murine and human type II IL-1 receptor proteins to their natural counterparts, and to natural and recombinant murine and human type I IL-1 receptors. In general, the sizes of the transiently-expressed recombinant receptors are similar to the natural receptors, although the recombinant proteins migrate slightly faster and as slightly broader bands, possibly as a result of differences in glycosylation pattern when over-expressed in CV1/EBNA cells. The results also indicate that the type II IL-1 receptors are smaller than the type I IL-1 receptors. One particular combination (natural human type I receptor with IL-1 $\beta$ ) failed to yield specific crosslinking products. Since approximately equal amounts of label were loaded into each experimental lane, as indicated by the intensity of the free ligand bands at the bottom of the gels, this combination must crosslink relatively poorly.

The lane showing natural human type II IL-1 receptor-bearing cells cross-linked with  $^{125}\text{I}$ -IL-1 $\alpha$  reveals a component in the size range ( $M_r=100,000$ ) of complexes with natural and recombinant type I receptors. No such complex can be detected in the lane containing recombinant type II IL-1 receptor, possibly as a result of low level expression of type I IL-1 receptors on the CB23 cells, since these cells contain trace amounts of type I IL-1 receptor mRNA.

### Example 7

#### Preparation of Monoclonal Antibodies to Type II IL-1R

Preparations of purified recombinant type II IL-1R, for example, human type II IL-1R, or transfected COS cells expressing high levels of type II IL-1R are employed to generate monoclonal antibodies against type II IL-1R using conventional techniques, for example, those disclosed in U.S. Patent 4,411,993. Such antibodies are likely to be useful in interfering with IL-1 binding to type II IL-1R, for example, in ameliorating toxic or other undesired effects of IL-1, or as components of diagnostic or research assays for IL-1 or soluble type II IL-1R.

To immunize mice, type II IL-1R immunogen is emulsified in complete Freund's adjuvant and injected in amounts ranging from 10-100 µg subcutaneously and intraperitoneally into Balb/c mice. Ten to twelve days later, the immunized animals are

boosted with additional immunogen emulsified in incomplete Freund's adjuvant and periodically boosted thereafter on a weekly to biweekly immunization schedule. Serum samples are periodically taken by retro-orbital bleeding or tail-tip excision for testing by dot-blot assay (antibody sandwich) or ELISA (enzyme-linked immunosorbent assay), or receptor binding inhibition. Other assay procedures are also suitable. Following detection of an appropriate antibody titer, positive animals are given an intravenous injection of antigen in saline. Three to four days later, the animals are sacrificed, splenocytes harvested, and fused to the murine myeloma cell line NS1 or Ag8.653. Hybridoma cell lines generated by this procedure are plated in multiple microtiter plates in a HAT selective medium (hypoxanthine, aminopterin, and thymidine) to inhibit proliferation of non-fused cells, myeloma hybrids, and spleen cell hybrids.

Hybridoma clones thus generated can be screened by ELISA for reactivity with type II IL-1R, for example, by adaptations of the techniques disclosed by Engvall et al., *Immunochem.* 8:871 (1971) and in U.S. Patent 4,703,004. Positive clones are then injected into the peritoneal cavities of syngeneic Balb/c mice to produce ascites containing high concentrations (>1 mg/ml) of anti-type II IL-1R monoclonal antibody, or grown in flasks or roller bottles. The resulting monoclonal antibody can be purified by ammonium sulfate precipitation followed by gel exclusion chromatography, and/or affinity chromatography based on binding of antibody to Protein A of *Staphylococcus aureus* or protein G from Streptococci.

#### **BRIEF DESCRIPTION OF THE SEQUENCE LISTING**

SEQ ID NO:1 and SEQ ID NO:2 show the nucleotide sequence and predicted amino acid sequence of human type II IL-1R. The mature peptide encoded by this sequence is defined by amino acids 1-385. The predicted signal peptide is defined by amino acids -13 through -1. The predicted transmembrane region is defined by amino acids 331-356.

SEQ ID NO:3 - SEQ ID NO:6 are various oligonucleotides used to clone the full-length human type II IL-1R.

SEQ ID NO:7 and SEQ ID NO:8 are oligonucleotide primers used to construct a soluble human type II IL-1R by polymerase chain reaction (PCR).

SEQ ID NO:9 - SEQ ID NO:11 are oligonucleotide primers used to clone a full-length and soluble murine type II IL-1Rs.

SEQ ID NO:12 and SEQ ID NO:13 show the nucleotide sequence and predicted amino acid sequence of the full-length murine type II IL-1R. The mature peptide encoded by this sequence is defined by amino acids 1-397. The predicted signal peptide is defined by amino acids -13 through -1. The predicted transmembrane region is defined by amino acids 343-368.

SEQ ID NO:14 is an oligonucleotide primer used to construct a soluble murine type II IL-1R.

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

- (i) APPLICANT: Sims, John E.  
Cosman, David J.  
Lupton, Stephen D.  
Mosley, Bruce A.  
Dower, Steven K.
- (ii) TITLE OF INVENTION: Type II Interleukin-1 Receptors
- (iii) NUMBER OF SEQUENCES: 14
- (iv) CORRESPONDENCE ADDRESS:
  - (A) ADDRESSEE: Immunex Corporation
  - (B) STREET: 51 University Street
  - (C) CITY: Seattle
  - (D) STATE: WA
  - (E) COUNTRY: USA
  - (F) ZIP: 98101
- (v) COMPUTER READABLE FORM:
  - (A) MEDIUM TYPE: Floppy disk
  - (B) COMPUTER: IBM PC Compatible
  - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
  - (D) SOFTWARE: PatentIn Release #1.24
- (vi) CURRENT APPLICATION DATA:
  - (A) APPLICATION NUMBER:
  - (B) FILING DATE: 16-MAY-1991
  - (C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
  - (A) APPLICATION NUMBER: US 07/534,193
  - (B) FILING DATE: 05-JUN-1990
- (viii) PRIOR APPLICATION DATA:
  - (A) APPLICATION NUMBER: US 07/573,576
  - (B) FILING DATE: 24-AUG-1990
- (ix) PRIOR APPLICATION DATA:
  - (A) APPLICATION NUMBER: 07/627,071
  - (B) FILING DATE: 13-DEC-1990
- (x) ATTORNEY/AGENT INFORMATION:
  - (A) NAME: Wight, Christopher L.
  - (B) REGISTRATION NUMBER: 31680
  - (C) REFERENCE/DOCKET NUMBER: 2003-C
- (xi) TELECOMMUNICATION INFORMATION:
  - (A) TELEPHONE: 206-587-5570
  - (B) TELEFAX: 206-233-0644

## (2) INFORMATION FOR SEQ ID NO:1:

- (1) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 1357 base pairs

(B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA to mRNA

(iii) HYPOTHETICAL: N

(iv) ANTI-SENSE: N

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Homo sapiens  
 (G) CELL TYPE: Human B cell lymphoblastoid  
 (H) CELL LINE: CB23

(vii) IMMEDIATE SOURCE:

(A) LIBRARY: CB23 cDNA  
 (B) CLONE: pHuIL-1RII75

(ix) FEATURE:

(A) NAME/KEY: CDS  
 (B) LOCATION: 154..1350  
 (D) OTHER INFORMATION:

(ix) FEATURE:

(A) NAME/KEY: mat\_peptide  
 (B) LOCATION: 193..1347  
 (D) OTHER INFORMATION:

(ix) FEATURE:

(A) NAME/KEY: sig\_peptide  
 (B) LOCATION: 154..192  
 (D) OTHER INFORMATION:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

CTGGAAAATA CATTCTGCTA CTCTTAAAAA CTAGTGACGC TCATACAAAT CAACAGAAAG	60
AGCTTCTGAA GGAAGACTTT AAAGCTGCTT CTGCCACGTG CTGCTGGGTC TCAGTCCTCC	120
ACTTCCCGTG TCCTCTGGAA GTTGTCAGGA GCA ATG TTG CGC TTG TAC GTG TTG	174
Met Leu Arg Leu Tyr Val Leu	
-13 -10	
GTA ATG GGA GTT TCT GCC TTC ACC CTT CAG CCT GCG GCA CAC ACA GGG	222
Val Met Gly Val Ser Ala Phe Thr Leu Gln Pro Ala Ala His Thr Gly	
-5 1 5 10	
GCT GCC AGA AGC TGC CGG TTT CGT GGG AGG CAT TAC AAG CGG GAG TTC	270
Ala Ala Arg Ser Cys Arg Phe Arg Gly Arg His Tyr Lys Arg Glu Phe	
15 20 25	
AGG CTG GAA GGG GAG CCT GTA GCC CTG AGG TGC CCC CAG GTG CCC TAC	318
Arg Leu Glu Gly Glu Pro Val Ala Leu Arg Cys Pro Gln Val Pro Tyr	
30 35 40	
TGG TTG TGG GCC TCT GTC AGC CCC CGC ATC AAC CTG ACA TGG CAT AAA	366
Trp Leu Trp Ala Ser Val Ser Pro Arg Ile Asn Leu Thr Trp His Lys	
45 50 55	

AAT GAC TCT GCT AGG ACG GTC CCA GGA GAA GAG ACA CGG ATG TGG Asn Asp Ser Ala Arg Thr Val Pro Gly Glu Glu Glu Thr Arg Met Trp 60 65 70	414
GCC CAG GAC GGT GCT CTG TGG CTT CTG CCA GCC TTG CAG GAG GAC TCT Ala Gln Asp Gly Ala Leu Trp Leu Leu Pro Ala Leu Gln Glu Asp Ser 75 80 85 90	462
GGC ACC TAC GTC TGC ACT ACT AGA AAT GCT TCT TAC TGT GAC AAA ATG Gly Thr Tyr Val Cys Thr Thr Arg Asn Ala Ser Tyr Cys Asp Lys Met 95 100 105	510
TCC ATT GAG CTC AGA GTT TTT GAG AAT ACA GAT GCT TTC CTG CCG TTC Ser Ile Glu Leu Arg Val Phe Glu Asn Thr Asp Ala Phe Leu Pro Phe 110 115 120	558
ATC TCA TAC CCG CAA ATT TTA ACC TTG TCA ACC TCT GGG GTA TTA GTA Ile Ser Tyr Pro Gln Ile Leu Thr Leu Ser Thr Ser Gly Val Leu Val 125 130 135	606
TGC CCT GAC CTG AGT GAA TTC ACC CGT GAC AAA ACT GAC GTG AAG ATT Cys Pro Asp Leu Ser Glu Phe Thr Arg Asp Lys Thr Asp Val Lys Ile 140 145 150	654
CAA TGG TAC AAG GAT TCT CTT CTT TTG GAT AAA GAC AAT GAG AAA TTT Gln Trp Tyr Lys Asp Ser Leu Leu Leu Asp Lys Asp Asn Glu Lys Phe 155 160 165 170	702
CTA AGT GTG AGG GGG ACC ACT CAC TTA CTC GTA CAC GAT GTG GCC CTG Leu Ser Val Arg Gly Thr Thr His Leu Leu Val His Asp Val Ala Leu 175 180 185	750
GAA GAT GCT GGC TAT TAC CGC TGT GTC CTG ACA TTT GCC CAT GAA GGC Glu Asp Ala Gly Tyr Tyr Arg Cys Val Leu Thr Phe Ala His Glu Gly 190 195 200	798
CAG CAA TAC AAC ATC ACT AGG AGT ATT GAG CTA CGC ATC AAG AAA AAA Gln Gln Tyr Asn Ile Thr Arg Ser Ile Glu Leu Arg Ile Lys Lys Lys 205 210 215	846
AAA GAA GAG ACC ATT CCT GTG ATC ATT TCC CCC CTC AAG ACC ATA TCA Lys Glu Glu Thr Ile Pro Val Ile Ile Ser Pro Leu Lys Thr Ile Ser 220 225 230	894
GCT TCT CTG GGG TCA AGA CTG ACA ATC CCG TGT AAG GTG TTT CTG GGA Ala Ser Leu Gly Ser Arg Leu Thr Ile Pro Cys Lys Val Phe Leu Gly 235 240 245 250	942
ACC GGC ACA CCC TTA ACC ACC ATG CTG TGG TGG ACG GCC AAT GAC ACC Thr Gly Thr Pro Leu Thr Thr Met Leu Trp Trp Thr Ala Asn Asp Thr 255 260 265	990
CAC ATA GAG AGC GCC TAC CCG GGA GGC CGC GTG ACC GAG GGG CCA CGC His Ile Glu Ser Ala Tyr Pro Gly Gly Arg Val Thr Glu Gly Pro Arg 270 275 280	1038
CAG GAA TAT TCA GAA AAT AAT GAG AAC TAC ATT GAA GTG CCA TTG ATT Gln Glu Tyr Ser Glu Asn Asn Glu Asn Tyr Ile Glu Val Pro Leu Ile 285 290 295	1086

TTT GAT CCT GTC ACA AGA GAG GAT TTG CAC ATG GAT TTT AAA TGT GTT 1134  
 Phe Asp Pro Val Thr Arg Glu Asp Leu His Met Asp Phe Lys Cys Val  
 300 305 310  
 GTC CAT AAT ACC CTG AGT TTT CAG ACA CTA CGC ACC ACA GTC AAG GAA 1182  
 Val His Asn Thr Leu Ser Phe Gln Thr Leu Arg Thr Thr Val Lys Glu  
 315 320 325 330  
 GCC TCC TCC ACG TTC TCC TGG GGC ATT GTG CTG GCC CCA CTT TCA CTG 1230  
 Ala Ser Ser Thr Phe Ser Trp Gly Ile Val Leu Ala Pro Leu Ser Leu  
 335 340 345  
 GCC TTC TTG GTT TTG GGG GGA ATA TGG ATG CAC AGA CGG TGC AAA CAC 1278  
 Ala Phe Leu Val Leu Gly Gly Ile Trp Met His Arg Arg Cys Lys His  
 350 355 360  
 AGA ACT GGA AAA GCA GAT GGT CTG ACT GTG CTA TGG CCT CAT CAT CAA 1326  
 Arg Thr Gly Lys Ala Asp Gly Leu Thr Val Leu Trp Pro His His Gln  
 365 370 375  
 GAC TTT CAA TCC TAT CCC AAG TGA AATAAAT 1357  
 Asp Phe Gln Ser Tyr Pro Lys  
 380 385

## (2) INFORMATION FOR SEQ ID NO:2:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 398 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met Leu Arg Leu Tyr Val Leu Val Met Gly Val Ser Ala Phe Thr Leu  
 -13 -10 -5 1  
 Gln Pro Ala Ala His Thr Gly Ala Ala Arg Ser Cys Arg Phe Arg Gly  
 5 10 15  
 Arg His Tyr Lys Arg Glu Phe Arg Leu Glu Gly Glu Pro Val Ala Leu  
 20 25 30 35  
 Arg Cys Pro Gln Val Pro Tyr Trp Leu Trp Ala Ser Val Ser Pro Arg  
 40 45 50  
 Ile Asn Leu Thr Trp His Lys Asn Asp Ser Ala Arg Thr Val Pro Gly  
 55 60 65  
 Glu Glu Glu Thr Arg Met Trp Ala Gln Asp Gly Ala Leu Trp Leu Leu  
 70 75 80  
 Pro Ala Leu Gln Glu Asp Ser Gly Thr Tyr Val Cys Thr Thr Arg Asn  
 85 90 95  
 Ala Ser Tyr Cys Asp Lys Met Ser Ile Glu Leu Arg Val Phe Glu Asn  
 100 105 110 115

39

Thr Asp Ala Phe Leu Pro Phe Ile Ser Tyr Pro Gln Ile Leu Thr Leu  
 120 125 130  
 Ser Thr Ser Gly Val Leu Val Cys Pro Asp Leu Ser Glu Phe Thr Arg  
 135 140 145  
 Asp Lys Thr Asp Val Lys Ile Gln Trp Tyr Lys Asp Ser Leu Leu Leu  
 150 155 160  
 Asp Lys Asp Asn Glu Lys Phe Leu Ser Val Arg Gly Thr Thr His Leu  
 165 170 175  
 Leu Val His Asp Val Ala Leu Glu Asp Ala Gly Tyr Tyr Arg Cys Val  
 180 185 190 195  
 Leu Thr Phe Ala His Glu Gly Gln Gln Tyr Asn Ile Thr Arg Ser Ile  
 200 205 210  
 Glu Leu Arg Ile Lys Lys Lys Lys Glu Glu Thr Ile Pro Val Ile Ile  
 215 220 225  
 Ser Pro Leu Lys Thr Ile Ser Ala Ser Leu Gly Ser Arg Leu Thr Ile  
 230 235 240  
 Pro Cys Lys Val Phe Leu Gly Thr Gly Thr Pro Leu Thr Thr Met Leu  
 245 250 255  
 Trp Trp Thr Ala Asn Asp Thr His Ile Glu Ser Ala Tyr Pro Gly Gly  
 260 265 270 275  
 Arg Val Thr Glu Gly Pro Arg Gln Glu Tyr Ser Glu Asn Asn Glu Asn  
 280 285 290  
 Tyr Ile Glu Val Pro Leu Ile Phe Asp Pro Val Thr Arg Glu Asp Leu  
 295 300 305  
 His Met Asp Phe Lys Cys Val Val His Asn Thr Leu Ser Phe Gln Thr  
 310 315 320  
 Leu Arg Thr Thr Val Lys Glu Ala Ser Ser Thr Phe Ser Trp Gly Ile  
 325 330 335  
 Val Leu Ala Pro Leu Ser Leu Ala Phe Leu Val Leu Gly Gly Ile Trp  
 340 345 350 355  
 Met His Arg Arg Cys Lys His Arg Thr Gly Lys Ala Asp Gly Leu Thr  
 360 365 370  
 Val Leu Trp Pro His His Gln Asp Phe Gln Ser Tyr Pro Lys  
 375 380 385

## (2) INFORMATION FOR SEQ ID NO:3:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA (genomic)

40

(iii) HYPOTHETICAL: N

(iv) ANTI-SENSE: N

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

TCGACTGGAA CGAGACGACC TGCT

24

2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 20 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: N

(iv) ANTI-SENSE: Y

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

GACCTTGCTC TGCTGGACGA

20

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 46 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: N

(iv) ANTI-SENSE: N

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

CTGTTGGGCT CGCGGTTGAG GACAACTCT TCGCGGTCTT TCCAGT 46

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 46 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: N

(iv) ANTI-SENSE: Y

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

GACAACCCGA GCGCCAACTC CTGTTTGAGA AGCGCCAGAA AGGTCA 46

(2) INFORMATION FOR SEQ ID NO:7:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 29 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: N

(iv) ANTI-SENSE: N

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

GCGTCGACCT AGTGACGCTC ATACAAATC 29

(2) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 33 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: N

(iv) ANTI-SENSE: N

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

GCGCGGCCGC TCAGGAGGAG GCTTCCTTGA CTG 33

(2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 37 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: N

(iv) ANTI-SENSE: N

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

CTGCAGGCGG CCGCGGATCC TTTTTTTTTT TTTTTT

37

(2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 28 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: N

(iv) ANTI-SENSE: N

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

GCGTCGACGG CAAGAAGCAG CAAGGTAC

28

(2) INFORMATION FOR SEQ ID NO:11:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 20 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: N

(iv) ANTI-SENSE: N

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

CTGCAGGCGG CCGCGGATCC

20

(2) INFORMATION FOR SEQ ID NO:12:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 1366 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA to mRNA

(iii) HYPOTHETICAL: N

(iv) ANTI-SENSE: N

(vi) ORIGINAL SOURCE:

(A) ORGANISM: Mouse

(H) CELL LINE: 70Z/3

## (vii) IMMEDIATE SOURCE:

(A) LIBRARY: 70Z/3

(B) CLONE: 12

## (ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 85..1317

(D) OTHER INFORMATION:

## (ix) FEATURE:

(A) NAME/KEY: mat\_peptide

(B) LOCATION: 124..1314

(D) OTHER INFORMATION:

## (ix) FEATURE:

(A) NAME/KEY: sig\_peptide

(B) LOCATION: 85..123

(D) OTHER INFORMATION:

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

GTCGACGGCA AGAAGCAGCA AGGTACAAGA ATACACAGCT CCAGGCTCCA AGGGTCCTGT	60
GCGCTCAGGA AGTTGGTGCG GACA ATG TTC ATC TTG CTT GTG TTA GTA ACT	111
Met Phe Ile Leu Leu Val Leu Val Thr	
-13 -10 -5	
GGA GTT TCT GCT TTC ACC ACT CCA ACA GTG GTG CAC ACA GGA AAG GTT	159
Gly Val Ser Ala Phe Thr Thr Pro Thr Val Val His Thr Gly Lys Val	
1 5 10	
TCT GAA TCC CCC ATT ACA TCG GAG AAG CCC ACA GTC CAT GGA GAC AAC	207
Ser Glu Ser Pro Ile Thr Ser Glu Lys Pro Thr Val His Gly Asp Asn	
15 20 25	
TGT CAG TTT CGT GGC AGA GAG TTC AAA TCT GAA TTG AGG CTG GAA GGT	255
Cys Gln Phe Arg Gly Arg Glu Phe Lys Ser Glu Leu Arg Leu Glu Gly	
30 35 40	
GAA CCT GTG GTT CTG AGG TGC CCC TTG GCA CCT CAC TCC GAC ATC TCC	303
Glu Pro Val Val Leu Arg Cys Pro Leu Ala Pro His Ser Asp Ile Ser	
45 50 55 60	
AGC AGT TCC CAT AGT TTT CTG ACC TGG AGT AAA TTG GAC TCT TCT CAG	351
Ser Ser Ser His Ser Phe Leu Thr Trp Ser Lys Leu Asp Ser Ser Gln	
65 70 75	
CTG ATC CCA AGA GAT GAG CCA AGG ATG TGG GTG AAG GGT AAC ATA CTC	399
Leu Ile Pro Arg Asp Glu Pro Arg Met Trp Val Lys Gly Asn Ile Leu	
80 85 90	
TGG ATT CTG CCA GCA GTG CAG CAA GAC TCT GGT ACC TAC ATT TGC ACA	447
Trp Ile Leu Pro Ala Val Gln Gln Asp Ser Gly Thr Tyr Ile Cys Thr	
95 100 105	
TTC AGA AAC GCA TCC CAC TGT GAG CAA ATG TCT GTG GAA CTC AAG GTC	495
Phe Arg Asn Ala Ser His Cys Glu Gln Met Ser Val Glu Leu Lys Val	
110 115 120	

44

TTT AAG AAT ACT GAA GCA TCT CTG CCT CAT GTC TCC TAC TTG CAA ATC Phe Lys Asn Thr Glu Ala Ser Leu Pro His Val Ser Tyr Leu Gln Ile 125 130 135 140	543
TCA GCT CTC TCC ACC ACC GGG TTA CTA GTG TGC CCT GAC CTG AAA GAA Ser Ala Leu Ser Thr Thr Gly Leu Leu Val Cys Pro Asp Leu Lys Glu 145 150 155	591
TTC ATC TCC AGC AAC GCT GAT GGA AAG ATA CAG TGG TAT AAG GGC GCC Phe Ile Ser Ser Asn Ala Asp Gly Lys Ile Gln Trp Tyr Lys Gly Ala 160 165 170	639
ATA CTC TTG GAT AAA GGC AAT AAG GAA TTT CTG AGT GCA GGA GAC CCC Ile Leu Leu Asp Lys Gly Asn Lys Glu Phe Leu Ser Ala Gly Asp Pro 175 180 185	687
ACA CGC CTA TTG ATA TCC AAC ACG TCC ATG GAC GAT GCA GGC TAT TAC Thr Arg Leu Leu Ile Ser Asn Thr Ser Met Asp Asp Ala Gly Tyr Tyr 190 195 200	735
AGA TGT GTT ATG ACA TTT ACC TAC AAT GGC CAG GAA TAC AAC ATC ACT Arg Cys Val Met Thr Phe Thr Tyr Asn Gly Gln Glu Tyr Asn Ile Thr 205 210 215 220	783
AGG AAT ATT GAA CTC CGG GTC AAA GGA GCA ACC ACG GAA CCC ATC CCT Arg Asn Ile Glu Leu Arg Val Lys Gly Ala Thr Thr Glu Pro Ile Pro 225 230 235	831
GTG ATC ATT TCT CCC CTG GAG ACA ATA CCA GCA TCA TTG GGG TCA AGA Val Ile Ile Ser Pro Leu Glu Thr Ile Pro Ala Ser Leu Gly Ser Arg 240 245 250	879
CTG ATA GTC CCG TGC AAA GTG TTT CTG GGA ACT GGT ACA TCT TCC AAC Leu Ile Val Pro Cys Lys Val Phe Leu Gly Thr Gly Thr Ser Ser Asn 255 260 265	927
ACC ATT GTG TGG TGG TTG GCT AAC AGC ACG TTT ATC TCG GCT GCT TAC Thr Ile Val Trp Trp Leu Ala Asn Ser Thr Phe Ile Ser Ala Ala Tyr 270 275 280	975
CCA AGA GGC CGT GTG ACC GAG GGG CTA CAC CAC CAG TAC TCA GAG AAT Pro Arg Gly Arg Val Thr Glu Gly Leu His His Gln Tyr Ser Glu Asn 285 290 295 300	1023
GAT GAA AAC TAT GTG GAA GTG TCG CTG ATT TTT GAT CCA GTC ACA AGG Asp Glu Asn Tyr Val Glu Val Ser Leu Ile Phe Asp Pro Val Thr Arg 305 310 315	1071
GAG GAT CTG CAT ACA GAT TTT AAA TGT GTT GCC TCG AAT CCA CGG AGT Glu Asp Leu His Thr Asp Phe Lys Cys Val Ala Ser Asn Pro Arg Ser 320 325 330	1119
TCT CAG TCA CTC CAT ACC ACA GTC AAA GAA GTC TCT TCC ACG TTC TCC Ser Gln Ser Leu His Thr Thr Val Lys Glu Val Ser Ser Thr Phe Ser 335 340 345	1167
TGG AGC ATT GCG CTG GCA CCT CTG TCT CTG ATC ATC TTG GTT GTG GGG Trp Ser Ile Ala Leu Ala Pro Leu Ser Leu Ile Ile Leu Val Val Gly 350 355 360	1215

GCA ATA TGG ATG CGC AGA CGG TGT AAA CGC AGG GCT GGA AAG ACA TAT	1263
Ala Ile Trp Met Arg Arg Arg Cys Lys Arg Arg Ala Gly Lys Thr Tyr	
365 370 375 380	
GGA CTG ACC AAG CTA CGG ACT GAC AAC CAG GAC TTC CCT TCC AGC CCA	1311
Gly Leu Thr Lys Leu Arg Thr Asp Asn Gln Asp Phe Pro Ser Ser Pro	
385 390 395	
AAC TAA ATAAAGGAAA TGAAATAAAA AAAAAAAAAA AAAAAGGATC CGCGGCCGC	1366
Asn .	

Met	Phe	Ile	Leu	Leu	Val	Leu	Val	Thr	Gly	Val	Ser	Ala	Phe	Thr	Thr
-13			-10					-5					1		
Pro	Thr	Val	Val	His	Thr	Gly	Lys	Val	Ser	Glu	Ser	Pro	Ile	Thr	Ser
	5					10					15				
Glu	Lys	Pro	Thr	Val	His	Gly	Asp	Asn	Cys	Gln	Phe	Arg	Gly	Arg	Glu
20					25					30					35
Phe	Lys	Ser	Glu	Leu	Arg	Leu	Glu	Gly	Glu	Pro	Val	Val	Leu	Arg	Cys
				40					45					50	
Pro	Leu	Ala	Pro	His	Ser	Asp	Ile	Ser	Ser	Ser	Ser	His	Ser	Phe	Leu
			55					60					65		
Thr	Trp	Ser	Lys	Leu	Asp	Ser	Ser	Gln	Leu	Ile	Pro	Arg	Asp	Glu	Pro
		70					75					80			
Arg	Met	Trp	Val	Lys	Gly	Asn	Ile	Leu	Trp	Ile	Leu	Pro	Ala	Val	Gln
	85					90					95				
Gln	Asp	Ser	Gly	Thr	Tyr	Ile	Cys	Thr	Phe	Arg	Asn	Ala	Ser	His	Cys
100					105					110					115
Glu	Gln	Met	Ser	Val	Glu	Leu	Lys	Val	Phe	Lys	Asn	Thr	Glu	Ala	Ser
				120					125					130	
Leu	Pro	His	Val	Ser	Tyr	Leu	Gln	Ile	Ser	Ala	Leu	Ser	Thr	Thr	Gly
			135					140					145		
Leu	Leu	Val	Cys	Pro	Asp	Leu	Lys	Glu	Phe	Ile	Ser	Ser	Asn	Ala	Asp
		150					155					160			
Gly	Lys	Ile	Gln	Trp	Tyr	Lys	Gly	Ala	Ile	Leu	Leu	Asp	Lys	Gly	Asn
	165					170					175				

46

Lys Glu Phe Leu Ser Ala Gly Asp Pro Thr Arg Leu Leu Ile Ser Asn  
 180 185 190 195  
 Thr Ser Met Asp Asp Ala Gly Tyr Tyr Arg Cys Val Met Thr Phe Thr  
 200 205 210  
 Tyr Asn Gly Gln Glu Tyr Asn Ile Thr Arg Asn Ile Glu Leu Arg Val  
 215 220 225  
 Lys Gly Ala Thr Thr Glu Pro Ile Pro Val Ile Ile Ser Pro Leu Glu  
 230 235 240  
 Thr Ile Pro Ala Ser Leu Gly Ser Arg Leu Ile Val Pro Cys Lys Val  
 245 250 255  
 Phe Leu Gly Thr Gly Thr Ser Ser Asn Thr Ile Val Trp Trp Leu Ala  
 260 265 270 275  
 Asn Ser Thr Phe Ile Ser Ala Ala Tyr Pro Arg Gly Arg Val Thr Glu  
 280 285 290  
 Gly Leu His His Gln Tyr Ser Glu Asn Asp Glu Asn Tyr Val Glu Val  
 295 300 305  
 Ser Leu Ile Phe Asp Pro Val Thr Arg Glu Asp Leu His Thr Asp Phe  
 310 315 320  
 Lys Cys Val Ala Ser Asn Pro Arg Ser Ser Gln Ser Leu His Thr Thr  
 325 330 335  
 Val Lys Glu Val Ser Ser Thr Phe Ser Trp Ser Ile Ala Leu Ala Pro  
 340 345 350 355  
 Leu Ser Leu Ile Ile Leu Val Val Gly Ala Ile Trp Met Arg Arg Arg  
 360 365 370  
 Cys Lys Arg Arg Ala Gly Lys Thr Tyr Gly Leu Thr Lys Leu Arg Thr  
 375 380 385  
 Asp Asn Gln Asp Phe Pro Ser Ser Pro Asn  
 390 395

## (2) INFORMATION FOR SEQ ID NO:14:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 36 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: DNA (genomic)

## (iii) HYPOTHETICAL: N

## (iv) ANTI-SENSE: N

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

GCGCGGCCGC CTAGGAAGAG ACTTCTTTGA CTGTGG

36

2501-WO

**CLAIMS**

1. An isolated DNA sequence encoding a biologically active type II IL-1 receptor (type II IL-1R) protein.
2. An isolated DNA sequence according to claim 1, selected from the group consisting of:
  - (a) cDNA clones having a nucleotide sequence derived from the coding region of a native type II IL-1R gene;
  - (b) DNA sequences capable of hybridization to the clones of (a) under moderately stringent conditions (50°C, 2 x SSC) and which encode biologically active type II IL-1R protein; and
  - (c) DNA sequences which are degenerate as a result of the genetic code to the DNA sequences defined in (a) and (b) and which encode biologically active type II IL-1R protein.
3. An isolated DNA sequence according to claim 1 which encodes a soluble type II IL-1R protein.
4. An isolated DNA sequence according to claim 3, wherein the soluble type II IL-1R protein is selected from the group consisting of the sequence of amino acid residues 1-333 of SEQ ID NO: 1 and the sequence of amino acid residues 1-345 of SEQ ID NO:12.
5. A recombinant expression vector comprising a DNA sequence according to any one of claims 1-4.
6. A process for preparing a biologically active mammalian type II IL-1 receptor (type II IL-1R) protein, comprising culturing a suitable host cell comprising a vector according to claim 5 under conditions promoting expression.
7. A purified biologically active type II IL-1 receptor (type II IL-1R) protein.
8. A purified biologically active soluble human type II IL-1R protein.
9. A purified biologically active type II IL-1R protein according to claim 7, selected from the group consisting of the sequence of amino acid residues 1-333 of SEQ ID NO: 1 and the sequence of amino acid residues 1-345 of SEQ ID NO:12.

10. A composition for regulating immune responses in a mammal, comprising an effective amount of a type II IL-1R protein composition according to claim 7, and a suitable diluent or carrier.

11. A method for regulating immune responses in a mammal, comprising administering an effective amount of a composition according to claim 10.

12. The method of claim 11, wherein the type II IL-1R protein is human type II IL-1R and the mammal to be treated is a human.

13. The use of a mammalian type II IL-1R protein in preparing a medicament for regulating immune responses in mammals.

14. The method of claim 13, wherein the type II IL-1R protein is human type II IL-1R and the mammal to be treated is a human.

15. The use of mammalian type II IL-1R protein in preparing a pharmaceutical composition suitable for parenteral administration to a human patient for regulating immune responses.

16. A process for detecting type II IL-1 or type II IL-1R molecules or the interaction thereof, comprising the steps of

- (a) binding a mammalian type II IL-1 receptor protein to an IL-1 molecule;
- and
- (b) detecting or measuring the amount of unbound IL-1 or type II IL-1 receptor.

17. Antibodies immunoreactive with and having specificity for mammalian type II IL-1 receptors.

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on 30/09/91. The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A- 8904838	01-06-89	US-A- 4968607	06-11-90
		AU-A- 2901589	14-06-89
		EP-A- 0318296	31-05-89
		JP-A- 2002351	08-01-90
WO-A- 9100742	24-01-91	AU-A- 6030990	06-02-91

## FURTHER INFORMATION CONTINUED FROM THE SECOND SHEET

V. ☒ OBSERVATION WHERE CERTAIN CLAIMS WERE FOUND UNSEARCHABLE <sup>1</sup>

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

- 1 ☒ Claim numbers 11,12 because they relate to subject matter not required to be searched by this Authority, namely

See PCT Rule 39.1(iv)

Methods for treatment of human or animal body by means of surgery or therapy, as well as diagnostic methods.

- 2 ☐ Claim numbers because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically

- 3 ☐ Claim numbers because they are dependent claims and are not drafted in accordance with the second and third sentences of PCT Rule 6(4a)

VI. ☐ OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING <sup>2</sup>

This international Searching Authority found multiple inventions in this international application as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims of the international application
2. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims of the international application for which fees were paid, specifically claims
3. ☐ No required additional search fees were timely paid by the applicant. Consequently this international search report is restricted to the invention first mentioned in the claims, it is covered by claim numbers
4. ☐ As all searchable claims could be searched without effort justifying an additional fee, the international Searching Authority did not invite payment of any additional fee

## Remark on Protest

- ☐ The additional search fees were accompanied by applicant's protest
- ☐ No protest accompanied the payment of additional search fees

## INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 91/03498

<b>I. CLASSIFICATION OF SUBJECT MATTER</b> (If several classification symbols apply, indicate all) <sup>9</sup>		
According to International Patent Classification (IPC) or to both National Classification and IPC		
Int.C1.5	C 12 N 15/12	C 07 K 13/00 C 12 P 21/08
G 01 N 33/68	A 61 K 37/02	
<b>II. FIELDS SEARCHED</b>		
Minimum Documentation Searched <sup>7</sup>		
Classification System	Classification Symbols	
Int.C1.5	C 12 N	C 07 K
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched <sup>8</sup>		
<b>III. DOCUMENTS CONSIDERED TO BE RELEVANT<sup>9</sup></b>		
Category <sup>10</sup>	Citation of Document, <sup>11</sup> with indication, where appropriate, of the relevant passages <sup>12</sup>	Relevant to Claim No. <sup>13</sup>
Y	FEBS LETTERS, vol. 243, no. 2, January 1989, G. SCAPIGLIATI et al.: "Differential binding of IL-1alpha and IL-1beta to receptors on B and T cells", pages 394-398, see the whole article, especially figure 5 ---	1-10,13 -17
Y	WO.A,8904838 (IMMUNEX CORP.) 1 June 1989, see the whole document ---	1-10,13 -17
Y	PROC. NATL. ACAD. SCI. USA, vol. 86, October 1989, K. BOMTZKY et al.: "Evidence for different interleukin 1 receptors in murine-B- and T-cell lines", pages 8034-8038, see the whole article, especially figure 2 (cited in the application) --- -/-	1-7,9, 10,13, 15-17
<p>* Special categories of cited documents:<sup>10</sup></p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"A" document member of the same patent family</p>		
<b>IV. CERTIFICATION</b>		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
11-09-1991	21 10 91	
International Searching Authority	Signature of Authorized Officer	
EUROPEAN PATENT OFFICE	M. PEIS <i>M. Peis</i>	

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No.
P,X	THE JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 265, no. 17, 15 June 1990, D. BENJAMIN et al.: "Heterogeneity in interleukin-1 (IL-1) receptors expressed on human B cell lines", pages 9943-9951, see the whole article ---	1-10,13 -17
P,X	WO,A,9100742 (DU PONT DE NEMOURS) 24 January 1991, see the whole document -----	1-10,13 -17

1/4




FIG. 1




FIG. 2


[illegible]

1	USSTFSUSIALAPLSLILUUGAIUM	ARACKRACKTYGLTKLATONQDPPSSPH*
2	ASSTESUGIUIAPISLAFULUGIUM	HAACKHATOKADOLTULUPHHQDQSYPK*

411  
398

**FIG. 3**

# Comparison of Natural and Recombinant IL-1 Receptors by Crosslinking

**FIG. 4**